INDEX FOR VOLUME 63

Compiled by Melissa R. Schneider

A

Abies amabilis, see Fir, Pacific silver
Abies balsamea, see Fir, balsam
Abies concolor, see Fir, white
Abies grandis, see Fir, grand
Abies lasiocarpa, see Fir, subalpine
Abraham, Kenneth F., James O. Leafloor, and Donald H. Rusch. Molt migrant Canada geese in northern Ontario and western James Bay, 649–655
Acacia berlandieri, see Giajillo
Acacia, blackbrush: 570
Acacia rigidula, see Acacia, blackbrush
Acacia smallii, see Huisache
Accipiter cooperii, see Hawk, Cooper's
Accipiter gentilis, see Goshawk
Accipiter nisus, see Hawk, sparrow
Accipiter striatus, see Hawk, sharp-shinned
Acer circinatum, see Maple, vine
Acer macrophyllum, see Maple, big-leaf
Acer negundo, see Boxelder
Acer rubrum, see Maple, red
Acer saccharinum, see Maple, silver
Acer saccharum, see Maple, sugar
Acer spicatum, see Maple, mountain
Acinonyx jubatus, see Cheetah
Acrididae, see Grasshopper
Acridotheres tristis, see Myna, common
Actitis macularia, see Sandpiper, spotted
Adams, Michael J. Correlated factors in amphibian decline: exotic species and habitat change in western Washington, 1162–1171
Adaptive kernel home range: 593–605, 1097
  effects of sample size on kernel home range estimates, 739–747
Adaptive resource management:
  effective monitoring for adaptive wildlife management: lessons from the Galápagos Islands, 1055–1065
  evaluation of waterfowl conservation under the North American Waterfowl Management Plan, 417–440
ADC, *see* Animal Damage Control
Africa: 1–13, 945
Afton, Alan D., *see* Pace, Richard M.
Agassiz National Wildlife Refuge: 389–395
Agave: 456
Agave spp, *see* Agave
Agelaius phoeniceus, see Blackbird, red-winged
Agouti: 3
Agouti paca, see Paca
Agriculture: 424–440, 519, 976–989, 1003–1008
  a regional model of the eastern cottontail and land use changes in Illinois, 956–963
  survival of ring-necked pheasant hens during spring in relation to landscape features, 147–154
Agropyron intermedium, see Grass, wheat
Agropyron smithii, see Grass, western wheat
Agropyron spicatum, see Grass, blue-bunch wheat
Agrostis alba, see Grass, redtop bent
AIC, *see* Akaike's Information Criterion
Aircraft:
  biases in bird strike statistics based on pilot reports, 997–1003
  response of fall-staging brant and Canada geese to aircraft overflights in southwestern Alaska, 373–381
Aix sponsa, see Duck, wood
Akaike's Information Criterion: 1111, 1121, 1241, 1243, 1254
Alabama: 26–35, 202, 223–231, 736
Alamosa National Wildlife Refuge: 129, 135
Alaska: 71, 104, 289, 335–345, 346, 373–381, 588–593, 662, 921–929, 1091, 1105, 1239, 1349–1358, 1378–1379
Alberta: 911–920, 1094–1108
Alces alces, see Moose
Alder, green: 168, 1307, 1321
Alder, red: 45, 485–486
Alder, smooth: 853–858, 1307
Alder, speckled: 575
Alectoris rufa, see Partridge
Alfalfa: 976, 981
Allantoin:
  sampling and statistical analysis of snow-urine allantoin:cretinine ratios, 1118–1132
Alligator, American: 890, 893
  effects of egg and hatchling harvest on American alligators in Florida, 1193–1200
Alligator mississipiansis, see Alligator, American
Allozyme variation:
  further evaluation of the genetic consequences of translocations on southeastern white-tailed deer population, 327–334
Alnus crispa, see Alder, green
Alnus rubra, see Alder, red
Alnus rugosa, see Alder, speckled
Alnus serrulata, see Alder, smooth
Alouatta pigra, see Monkey, black howler
Alpe, Michael J., James L. Kingery, and Jeffrey C. Mosley. Effects of summer sheep grazing on browse nutritive quality in autumn and winter, 346–354
Alpizar-Jara, Russell, *see* Pack, James C.
Ambrose, Skip, *see* Britten, Michael W.

INDEX FOR VOLUME 63

Compiled by Melissa R. Schneider

A

Abies amabilis, see Fir, Pacific silver
Abies balsamea, see Fir, balsam
Abies concolor, see Fir, white
Abies grandis, see Fir, grand
Abies lasiocarpa, see Fir, subalpine
Abraham, Kenneth F., James O. Leafloor, and Donald H. Rusch. Molt migrant Canada geese in northern Ontario and western James Bay, 649–655
Acacia berlandieri, see Giajillo
Acacia, blackbrush: 570
Acacia rigidula, see Acacia, blackbrush
Acacia smallii, see Huisache
Accipiter cooperii, see Hawk, Cooper's
Accipiter gentilis, see Goshawk
Accipiter nisus, see Hawk, sparrow
Accipiter striatus, see Hawk, sharp-shinned
Acer circinatum, see Maple, vine
Acer macrophyllum, see Maple, big-leaf
Acer negundo, see Boxelder
Acer rubrum, see Maple, red
Acer saccharinum, see Maple, silver
Acer saccharum, see Maple, sugar
Acer spicatum, see Maple, mountain
Acinonyx jubatus, see Cheetah
Acrididae, see Grasshopper
Acridotheres tristis, see Myna, common
Actitis macularia, see Sandpiper, spotted
Adams, Michael J. Correlated factors in amphibian decline: exotic species and habitat change in western Washington, 1162–1171
Adaptive kernel home range: 593–605, 1097
  effects of sample size on kernel home range estimates, 739–747
Adaptive resource management:
  effective monitoring for adaptive wildlife management: lessons from the Galápagos Islands, 1055–1065
  evaluation of waterfowl conservation under the North American Waterfowl Management Plan, 417–440
ADC, *see* Animal Damage Control
Africa: 1–13, 945
Afton, Alan D., *see* Pace, Richard M.
Agassiz National Wildlife Refuge: 389–395
Agave: 456
Agave spp, *see* Agave
Agelaius phoeniceus, see Blackbird, red-winged
Agouti: 3
Agouti paca, see Paca
Agriculture: 424–440, 519, 976–989, 1003–1008
  a regional model of the eastern cottontail and land use changes in Illinois, 956–963
  survival of ring-necked pheasant hens during spring in relation to landscape features, 147–154
Agropyron intermedium, see Grass, wheat
Agropyron smithii, see Grass, western wheat
Agropyron spicatum, see Grass, blue-bunch wheat
Agrostis alba, see Grass, redtop bent
AIC, *see* Akaike's Information Criterion
Aircraft:
  biases in bird strike statistics based on pilot reports, 997–1003
  response of fall-staging brant and Canada geese to aircraft overflights in southwestern Alaska, 373–381
Aix sponsa, see Duck, wood
Akaike's Information Criterion: 1111, 1121, 1241, 1243, 1254
Alabama: 26–35, 202, 223–231, 736
Alamosa National Wildlife Refuge: 129, 135
Alaska: 71, 104, 289, 335–345, 346, 373–381, 588–593, 662, 921–929, 1091, 1105, 1239, 1349–1358, 1378–1379
Alberta: 911–920, 1094–1108
Alces alces, see Moose
Alder, green: 168, 1307, 1321
Alder, red: 45, 485–486
Alder, smooth: 853–858, 1307
Alder, speckled: 575
Alectoris rufa, see Partridge
Alfalfa: 976, 981
Allantoin:
  sampling and statistical analysis of snow-urine allantoin:cretinine ratios, 1118–1132
Alligator, American: 890, 893
  effects of egg and hatchling harvest on American alligators in Florida, 1193–1200
Alligator mississipiansis, see Alligator, American
Allozyme variation:
  further evaluation of the genetic consequences of translocations on southeastern white-tailed deer population, 327–334
Alnus crispa, see Alder, green
Alnus rubra, see Alder, red
Alnus rugosa, see Alder, speckled
Alnus serrulata, see Alder, smooth
Alouatta pigra, see Monkey, black howler
Alpe, Michael J., James L. Kingery, and Jeffrey C. Mosley. Effects of summer sheep grazing on browse nutritive quality in autumn and winter, 346–354
Alpizar-Jara, Russell, *see* Pack, James C.
Ambrose, Skip, *see* Britten, Michael W.

Ambrosia dumosa, see Bursage
Ambystoma maculatum, see Salamander, spotted
Ambystoma spp., *see* Salamander, mole
Amelanchier alnifolia, see Serviceberry
Ammodramus savannarum, see Sparrow, grasshopper
Ammospermophilus nelsoni, see Squirrel, San Joaquin antelope
Amphibians: 484
correlated factors in amphibian decline: exotic species and habitat change in western Washington, 1162–1171
forest canopy closure and juvenile emigration by pool-breeding amphibians in Maine, 441–450
Amphispiza belli, see Sparrow, sage
Anabat II: 553–561, 1308–1320
Anas acuta, see Pintail, northern
Anas americana, see Wigeon, American
Anas clypeata, see Shoveler, northern
Anas cyanoptera, see Teal, cinnamon
Anas discors, see Teal, blue-winged
Anas fulvigula, see Duck, mottled
Anas rubripes, see Duck, American black
Anas strepera, see Gadwall
Anas wyvilliana, see Duck, Hawaiian
Andelt, William F., *see* Baker, Dan L.
Anderson, Gary B., *see* Stoops, Monica A.
Anderson, Stanley H., *see* Ayers, Loren W.
Andropogon gerardii, see Grass, big bluestem
Andropogon spp., *see* Grass, bluestem
Animal Damage Control: 278–286, 593–612, 997–1017, 1138–1151, 1327–1349
Anser albifrons frontalis, see Goose, greater white-fronted
Anser anser flavirostris, see Goose, Greenland white-fronted
Anser brachyrhynchus, see Goose, pink-footed
Ant, carpenter: 462, 467
Ant, red fire: 299
northern bobwhite chick mortality caused by red imported fire ants, 1291–1298
Antelope, pronghorn: 354, 923
Antennaria neglecta, see Everlasting
Anthony, Robert G, Monte G. Garrett, Frank B. Isaacs. Double-survey estimates of bald eagle populations in Oregon, 794–802
Anthony, Robert G., *see* Martin, Karl J.
Anthraquinone:
plant growth regulator (Stronghold™) enhances repellency of anthraquinone formulation (Flight Control™) to Canada geese, 1336–1343
Antilocarpa americana, see Antelope, pronghorn
ANWR, *see* Arctic National Wildlife Refuge
Aphelcoma coerulescens, see Jay, Florida scrub
Apple, crab: 309–310
Applejohn, Andrew P., *see* Jung, Thomas S.
Aquila chrysaetos, see Eagle, golden
Aquila nipalensis, see Eagle, steppe
Arbutus menziesii, see Madrone
Archilochus alexandri, see Hummingbird, black-chinned
Archilochus colubris, see Hummingbird, ruby-throated
Arctic National Wildlife Refuge: 588–593
Ardea herodias, see Heron, great blue
Arenaria interpres, see Turnstone, ruddy
Arengo, Felicity, and Guy A. Baldassarre. Resource variability and conservation of American flamingos in coastal wetlands of Yucatán, Mexico, 1201–1212
Aresco, Matthew J., and Craig Guyer. Burrow abandonment by gopher tortoises in slash pine plantations of the Conecuh National Forest, 26–35
Argentina: 895–900
Aristida stricta, see Grass, wire
Aristida spp., *see* Three-awn
Arizona: 77–84, 104, 154–161, 451–461, 546, 921–929, 1162, 1327, 1359–1358
Arkansas: 220, 424, 470–483, 1281–1290
ARM, *see* Adaptive resource management
Armadillo: 891
Artemia spp., *see* Shrimp, brine
Artemia tridentata, see Sagebrush, big
Artemisia pedatifida, see Sage, birdsfoot
Artemisia spp., *see* Sagebrush
Artemisia tridentata wyomingensis, see Sagebrush, Wyoming big
Arthur, Stephen M., *see* Jacoby, Michael E.
Ash, black: 575
Ash, green: 630–636, 640
Ash, velvet: 452
Asia: 6, 8
Asio flammeus, see Owl, short-eared
Aspen, quaking: 37, 39, 181, 316, 425, 462–466, 581, 1307
conifer release alternatives increase aspen forage quality in northwestern Ontario, 1320–1326
effectiveness of Hot Sauce® and Deer Away® repellents for deterring elk browsing of aspen sprouts, 1327–1336
Aster, woody: 272
***Astrocaryum* spp.:** 547
Atchafalaya Delta Wildlife Management Area: 138
Athene cunicularia, see Owl, burrowing
Atherurus africanus: 6–7
Atriplex confertifolia, see Shadscale
Atriplex gardnerii, see Saltbush, Gardner
Auklet, Cassin's: 15
Auklet, rhinoceros: 15
Aune, Keith, *see* Roffe, Thomas J.
Australia: 298–299, 306, 1140, 1145, 1193
Autocorrelation:
autocorrelation of location estimates and the analysis of radiotraking data, 1039–1044
Avery, Michael L., John S. Humphrey, David G. Decker, and Arlene P. McGrane. Seed color avoidance by captive red-winged blackbirds and boat-tailed grackles, 1003–1008

Avian botulism:
environmental characteristics associated with the occurrence of avian botulism in wetlands of a northern California refuge, 358–368
water and sediment characterics associated with avian botulism outbreaks in wetland, 1249–1260
Avicennia germinans, see Mangrove, white
Avocet, American:
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
Ayers, Loren W., and Stanley H. Anderson. An aerial sightability model for estimating ferruginous hawk population size, 85–97
Aythya affinis, see Scaup, lesser
Aythya americana, see Redhead
Aythya crecca, see Teal, green-winged
Aythya ferina, see Pochard, common
Aythya fuligula, see Duck, tufted
Aythya marila, see Scaup, greater
Aythya valisineria, see Canvasback

B

Baccharis angustifolia, see Silverling
Baccharis glomeruliflora, see Tree, groundsel
***Bactris oligocarpa*:** 547
Badger: 270, 275, 1066–1081, 1300
Badlands National Park: 275
Baeolophus bicolor, see Titmouse, tufted
Baker, Bruce W., Thomas R. Stanley, and James A. Sedgwick. Predation of artificial ground nests on white-tailed prairie dog colonies, 270–277
Baker, Dan L., William F. Andelt, Kenneth P. Burnham, and Wayne D. Shepperd. Effectiveness of Hot Sauce® and Deer Away® repellents for deterring elk browsing of aspen sprouts, 1327–1336
Baldassarre, Guy A., *see* Arengo, Felicity
Ballard, Warren B., Heather A. Whitlaw, Steven J. Young, Roger A. Jenkins, and Graham J. Forbes. Predation and survival of white-tailed deer fawns in northcentral New Brunswick, 574–579
Bamboo: 233, 459
Bambus aculeata, see Bamboo
Bambusa vulgaris, see Bamboo
Band, leg: 163, 640, 781, 1240
evaluation of a plasticine-filled leg band for day-old ducklings, 656–663
Band, wing: 163
Banff National Park: 1094
Bartlett, John G., *see* Dettmers, Randy
Bartmann, Richard M., *see* Unsworth, James W.
Bass, smallmouth: 1166
Bat, big brown: 1307–1320
Bat, eastern big brown: 1315
Bat, eastern small-footed: 1307–1302
Bat, hoary:
habitat selection by forest bats in relation to mixed-wood stand types and structure in central Ontario, 1306–1319
Bat, little brown:
habitat selection by forest bats in relation to mixed-wood stand types and structure in central Ontario, 1306–1319
Bat, northern long-eared:
habitat selection by forest bats in relation to mixed-wood stand types and structure in central Ontario, 1306–1319
Bat, red: 1307–1320
Bat, silver-haired: 553–561
habitat selection by forest bats in relation to mixed-wood stand types and structure in central Ontario, 1306–1319
Bat, Townsend's big-eared: 553–561
Bay, California: 45, 56
Bay, red: 874
Bay, sweet: 874
Beadlily, queencup: 347
Bear, black: 286, 305, 396, 574–579, 582, 608, 902, 915, 1104
dynamics of cervids in a multipredator ecosystem, 1082–1093
trophic relations of brown and black bears in several North American ecosystems, 921–929 species-specific population
Bear, brown: 335–336, 338, 341, 343, 354–357, 396
trophic relations of brown and black bears in several North American ecosystems, 921–929
Bear, grizzly: 396, 902, 1104
evaluating nutritional condition of grizzly bears via select blood parameters, 286–281 rates and causes of grizzly bear mortality in the interior mountains of British Columbia, Alberta, Montana, Washington, and Idaho, 911–920
species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
Beault, André, *see* Lougheed, Lynn W.
Beaver: 313, 404, 443, 646, 854, 859, 1307
Beaver ponds: 424–425
Beech: 512
Beech, American: 181, 442, 462, 965
Beier, Paul, *see* Delany, David K.
Belize: 748–755
Below, Theodore H., *see* Mumme, Ronald L.
Berberis nervosa, see Grape, Oregon
Betula alleghaniensis, see Birch, yellow
Betula glandulosa, see Birch, dwarf
Betula nana, see Birch, dwarf
Betula papyrifera, see Birch, white
Betula pendula, see Birch, silver
Betula populifolia, see Birch, gray
Betula pubescens var. *tortuosa, see* Birch, valley
BIA, *see* Bioelectrical impedance analysis
Bias: 85–97, 1105, 1372
biases in bird strike statistics based on pilot reports, 997–1003
Big Cypress National Preserve: 872–879

Bioaccumulation: 922
selenium accumulation and effects on reproduction in captive American kestrels fed selenomethionine, 502–511
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
selenium in blood of predatory birds from Kesterson Reservoir and other areas in California, 1273–1281
Bioelectrical impedance analysis:
evaluating nutritional condition of grizzly bears via select blood parameters, 286–281
Biological significance: 1360
the insignificance of statistical significance testing, 763–772
Birch, dwarf: 287, 336, 581
Birch, gray: 443
Birch, silver: 355
Birch, valley: 728
Birch, white: 168, 181, 355, 462, 575, 581, 1307, 1321
Birch, yellow: 168, 181, 442, 462–466, 965
Bison:
biosafety of parenteral *Brucella abortus* RB51 vaccine in bison calves, 950–955
brucellosis in Yellowstone National Park bison: quantitative serology and infection, 1132–1137
Bison bison, see Bison
Bissonette, John A., *see* Stolley, Dorie S.
Bitterbush, antelope: 86, 316–317, 347
Blackbird, red-winged: 530–538, 1233
seed color avoidance by captive red-winged blackbirds and boat-tailed grackles, 1003–1008
Blackbird, rusty: 170–177
Blackwell, Bradley F., Thomas W. Seamans, and Richard A. Dolbeer. Plant growth regulator (Stronghold™) enhances repellency of anthraquinone formulation (Flight Control™) to Canada geese, 1336–1343
Blarina brevicauda, see Shrew, northern short-tailed
Blejwas, Karen M., *see* Sacks, Benjamin N.
Blood chemistry: 930–938
evaluating nutritional condition of grizzly bears via select blood parameters, 286–281
real-time physiological responses of red deer to translocations, 1152–1162
selenium in blood of predatory birds from Kesterson Reservoir and other areas in California, 1273–1281
Blueberry: 1307
Bluebird, eastern: 807, 1013
Blums, Peter, J. Brian Davis, Scott E. Stephens, Aivars Mednis, and David M. Richardson. Evaluation of a plasticine-filled leg band for day-old ducklings, 656–663
Boal, Clint W., and R. William Mannan. Comparative breeding ecology of Cooper's hawks in urban and exurban areas of southeastern Arizona, 77–84
Bobcat: 305, 574–579, 890, 907, 935, 1066–1081
influence of time-in-residence on home range and habitat use of bobcats, 261–269
survival and cause-specific mortality of adult bobcats in central Mississippi, 613–620
Bobolink: 532
Bodmer, Richard E., *see* Robinson, John G.
Body mass: 315–326, 346, 695–704, 930–938, 993–996, 1046–1050, 1128, 1138–1151, 1240–1248, 1282–1290
Bogenschutz, Todd R., *see* Clark, William R.
Boiga irregularis, see Snake, brown tree
Bombycilla cedrorum, see Waxwing, cedar
Bonasa umbellus, see Grouse, ruffed
Bone marrow: 946
Borkowski, John J., *see* Pils, Andrew C.
Bosque National Wildlife Refuge: 130
Bothriochloa ischaemum, see Grass, plains bluestem
Bouteloua curtipendula, see Grama, sideoats
Bouteloua eripoda, see Grama, black
Bouteloua gracilis, see Grama, blue
Boutin, Stan, *see* Marshal, Jason P.
Boxelder: 640
Boyce, Walter M., *see* Waithman, John D.
Boyd, Diane K., and Daniel H. Pletscher. Characteristics of dispersal in a colonizing wolf population in the central Rocky Mountains, 1094–1108
Brant, Pacific:
response of fall-staging brant and Canada geese to aircraft overflights in southwestern Alaska, 373–381
Branta bernicla nigricans, see Brant, Pacific
Branta canadensis canadensis, see Goose, Canada
Branta canadensis interior, see Goose, Canada
Branta canadensis maxima, see Goose, giant Canada
Branta canadensis minima, see Goose, cackling Canada
Branta canadensis moffitti, see Goose, western Canada
Branta canadensis, see Goose, Canada
Branta canadensis taverneri, see Goose, Canada
Branta leucopsis, see Goose, barnacle
Brasher, Michael G., *see* Gray, Matthew J.
Brazil: 1–13, 570
Breault, André *see* Lougheed, Lynn W.
Brennan, Leonard A., *see* Carver, A. Vincent
Brinkhaus, Amy J., *see* Waithman, John D.
British Columbia: 14–25, 346, 585, 784, 902, 911–920, 1094–1108, 1359–1369, 1383
Britten, Michael W., Patricia L. Kennedy, and Skip Ambrose. Performance and accuracy evaluation of small satellite transmitters, 1349–1358
Bro, Elisabeth, Jean Clobert, and François Reitz. Effects of radiotransmitters on survival and reproductive success of gray partridge, 1044–1051

Brome, fringed: 1328
Brome, smooth: 517, 687, 712, 976, 981
Bromus cilatus, see Brome, fringed
Bromus inermis, see Brome, smooth
Bromus tectorum, see Grass, cheat
Brooks, Robert T. Residual effects of thinning and high white-tailed deer densities on northern red-back salamanders in southern New England oak forests, 1172–1180
Brown, W. Kent, *see* Schaefer, James A.
Brown, Wendy M., *see* Drewien, Roderick C.
Brownie, Cavell, *see* Tsai, Kuenhi
Browse:
effects of summer sheep grazing on browse nutritive quality in autumn and winter, 346–354
Brucella abortus, see Brucellosis
Brucellosis: 927
biosafety of parenteral *Brucella abortus* RB51 vaccine in bison calves, 950–955
brucellosis in Yellowstone National Park bison: quantitative serology and infection, 1132–1137
Brundige, Gary C., *see* Seaman, D. Erran
Bryant, Fred C., *see* Henke, Scott E.
Bubo virginianus, see Owl, great-horned
Bucephala albeola, see Bufflehead
Bucephala islandica, see Goldeneye, Barrow's
Buchloe dactyloides, see Grass, buffalo
Buckley, P. A., *see* Legare, Michael L.
Buehler, David A., *see* Dettmers, Randy
Buffalo berry: 912
Buffer, riparian:
breeding bird assemblages inhabiting riparian buffer strips in Newfoundland, Canada, 167–179
influence of riparian buffer width on bird assemblages in western Oregon, 484–496
Bufflehead: 1363–1364
Bufo americanus, see Toad, American
Buford, Ernest W., and David E. Capen. Abundance and productivity of forest songbirds in a managed, unfragmented landscape in Vermont, 180–188
Bullhead: 1166
Bulrush: 640, 853–858
Bulrush, hardstem: 359
Bunting, indigo: 803–814, 817–821
Bunting, lark: 530–538
Bunting, painted: 807
Bureau of Land Management: 485, 1182, 1213
Burger, Loren W., Jr., *see* Carver, A. Vincent
———, *see* Frawley, Brian J.
———, *see* McCoy, Timothy D.
———, *see* Chamberlain, Michael J.
Burnham, Kenneth P., *see* Baker, Dan L.
Burrows:
burrow abandonment by gopher tortoises in slash pine plantations of the Conecuh National Forest, 26–35
norway rats as predators of burrow-nesting seabirds: insights from stable isotope analyses, 14–25
Bursage: 1182
Bursera simaruba, see Limbo, gumbo
Buteo jamaicensis, see Hawk, red-tailed
Buteo platypterus, see Hawk, broad-winged
Buteo regalis, see Hawk, ferruginous
Buteo swainsoni, see Hawk, Swainson's
Buttonwood: 251

C

Cacti: 456
Cactus, pencil cholla: 1183
Cactus, pricklypear: 571
Calamagrostis nuktaensis, see Grass, reed
Calathea dilabens: 547
Calathea zingiberina: 547
Calcarius mccownii, see Longspur, McCown's
Calcarius ornatus, see Longspur, chestnut-collared
California: 44–59, 292, 298–308, 358–368, 492, 497–501, 502–511, 561–569, 591, 593–605, 662, 784, 786–787, 930–949, 1079, 1162–1171, 1219, 1222–1238, 1239–1249, 1273–1281
Callamospiza melanocorys, see Bunting, lark
Callipepla gambelii, see Quail, Gambel's
Callipepla squamata, see Quail, scaled
Camponotus spp., *see* Ant, carpenter
Canada: 14–25, 167–179, 283, 287–291, 343, 382–389, 417–440, 574–579, 649–655, 662, 933, 1258
Cane, maiden: 874
Canine parvovirus: 930–938
Canis mesomelas, see Jackal, black-backed
Canis familiaris dingo, see Dingo
Canis familiaris, see Dog, domestic
Canis latrans, see Coyote
Canis lupus pallipes, see Wolf
Canis lupus, see Wolf, gray
Canis simensis, see Wolf, Ethiopian
Canopy closure: 26–35, 337, 339, 484, 492, 494
forest canopy closure and juvenile emigration by pool-breeding amphibians in Maine, 441–450
Canvasback: 393, 418, 429–430, 647, 1363–1364
Canyons: 451–461
Capen, David E., *see* Buford, Ernest W.
Capreolus capreolus, see Deer, roe
Caprimulgus, vociferus, see Whip-poor-will
Carbamates: 790
Carbon: 14–25
Cardinal, northern: 807–809, 1000
Cardinal, red-crested: 1000
Cardinalis cardinalis, see Cardinal, northern
Carduelis pinus, see Siskin, pine
Carduelis tristis, see Goldfinch, American
Carex geyeri, see Sedge, elk
Carex nesophila, see Sedge, short-stalk
Carex spp., *see* Sedge
Caribou: 289, 397, 567, 728, 1090, 1110, 1159
caribou calf production and seasonal range quality during a population decline, 335–345

use of climatic data to model the presence of oestrid flies in caribou herds, 588–593
Caribou National Forest: 607
Caribou, woodland:
demography of decline of the Red Wine Mountains caribou herd, 580–587
Carlile, Lawrence D., *see* Mitchell, Lee R.
Carney, Daniel W., *see* McLellan, Bruce N.
Carpenter, Leslie B., *see* Kochert, Michael N.
Carpodacus mexicanus, see Finch, house
Carpodacus purpureus, see Finch, purple
Carrie, Dawn K., *see* Carrie, N. Ross
Carrie, N. Ross, Richard N. Conner, D. Craig Rudolph, and Dawn K. Carrie. Reintroduction and postrelease movements of red-cockaded woodpecker groups in eastern Texas, 824–832
Carrying capacity:
energy-based carrying capacity for quails, 664–67
CART, *see* Classification and regression tree analysis
Carter, Shawn L., Carola A. Haas, and Joseph C. Mitchell. Home range and habitat selection of bog turtles in southwestern Virginia, 853–860
Carver, A. Vincent, Loren W. Burger, Jr., and Leonard A. Brennan. Passive integrated transponders and patagial tag markers for northern bobwhite chicks, 162–166
Cary, John R., *see* Taylor, J. Scott
Carya ovata, see Hickory, shagbark
Carya tomentosa, see Hickory, mockernut
Case, Ray, *see* Gau, Robert J.
Cassini, Marcelo Hernán, *see* Guichón, María Laura
Castanea dentata, see Chestnut, American
Castor canadensis, see Beaver
Cat, feral: 256–257
Catbird, gray: 807–808
Cathartes aura, see Vulture, turkey
Catharus fuscescens, see Veery
Catharus guttatus, see Thrush, hermit
Catharus ustulatus, see Thrush, Swainson's
Cattail: 359, 640, 711–722, 898, 978
Causton, Charlotte E., *see* Gibbs, James P.
Ceanothus, redstem: 346–354
Ceanothus sanguineus, see Ceanothus, redstem
Cecropia schreberiana, see Tree, trumpet
Cedar, eastern white: 575
Cedar, incense: 292
Cedar, western red: 16, 292, 347, 485, 554, 1213
Cederlund, Goran, *see* Swenson, Jon E.
Celtis pallida, see Granjeno
Cenizo: 570
Centrocercus urophasianus, see Grouse, sage
Cephalophus monticola, see Duiker, blue
***Cephalophus ogibyi*:** 6–7
Cephalophus spp., *see* Duiker spp.
Cephanamyia trompe, see Fly, nasal bot
Ceratoides lanata, see Winterfat
Ceratopogonids: 362
Cercocarpus montanus, see Mahogany, mountain
***Cercopithecus erythrois*:** 6–7
***Cercopithecus mitis*:** 6–7
***Cercopithecus nictitans*:** 6–7
***Cercopithecus pogonias*:** 6–7
***Cercopithecus preussi*:** 6–7
Cerrorhinca monocerata, see Auklet, rhinoceros
Certhia americana, see Creeper, brown
Cervus elaphus, see Deer, red
Cervus elaphus nannodes, see Elk, tule
Cervus elaphus nelsoni, see Elk, Rocky Mountain
Cervus elaphus roosevelti, see Elk, Roosevelt
Cervus nippon, see Deer, Japanese sika
Ceryle alcyon, see Kingfisher, belted
Chaetodipus hispidus, see Mouse, hispid pocket
Chamaea fasciata, see Tit, wren
Chamberlain, Michael J., Bruce D. Leopold, Loren W. Burger, Jr., Bruce W. Plowman, and L. Michael Conner. Survival and cause-specific mortality of adult bobcats in central Mississippi, 613–620
Chamberlain, Michael J., Kurt M. Hodges, Bruce D. Leopold, and Tim S. Wilson. Survival and cause-specific mortality of adult raccoons in central Mississippi, 880–888
Chandler, C. Ray, *see* Mitchell, Lee R.
Chaoborids: 362
Chapman, Kim A., *see* Merrill, Michael D.
Chara spp., *see* Grass, musk
Charadrius montanus, see Plover, mountain
Charadrius vociferus, see Killdeer
Chat, Yellow-breasted: 806–808
Cheetah: 618, 906
Chen caerulescens atlantica, see Goose, greater snow
Chen caerulescens caerulescens, see Goose, lesser snow
Chen canagica, see Goose, emperor
Chen rossii, see Goose, Ross's
Cherokee National Forest: 816–823
Cherry, black: 965
Cherry, Marilyn L., *see* Lautenschlager, R. A.
Cherry, pin: 168, 1307
Chestnut, American: 965
Chickadee, black-capped: 170–177, 187–188, 489, 1017
Chickadee, boreal: 170–177
Chickadee, Carolina: 807
Chickadee, chestnut-backed: 484–496
Chicken, greater prairie: 517, 986
land-use patterns surrounding greater prairie-chicken leks in northwestern Minnesota, 189–198
Chihuahuan Desert: 1075
Chimpanzee: 753
Chironomids: 362
Chloris cucullata, see Grass, hooded windmill
Choristoneura fumiferana, see Spruce budworm
Christie, Jeffery S., *see* Krementz, David G.
Chrysobalanus icaco, see Cocoplum
Chrysothamnus spp., *see* Rabbitbrush
Church, Kevin E., *see* Taylor, J. Scott
Cinquefoil, marsh: 336

Circus cyaneus, see Harrier, northern
Cirsium spp., *see* Thistle
Cladina spp., *see* Lichen
Cladium jamaicensis, see Grass, saw
Clark, William R., Richard A. Schmitz, and Todd R. Bogenschutz. Site selection and nest success of ring-necked pheasants as a function of location in Iowa landscapes, 976–989
Clark, William R., *see* Schmitz, Richard A.
Classification and regression tree analysis: 956–963, 976–989
Clearcut: 167–179, 441–442, 486, 492, 812, 850, 1172–1180, 1386
Clemmys muhlengergii, see Turtle, bog
Clethrionomys gapperi, see Vole, southern red-backed
Clethrionomys rutilus, see Vole, northern red-backed
Climate:
effects of drought on desert tortoise movement and activity, 1181–1192
use of climatic data to model the presence of oestrid flies in caribou herds, 588–593
Clintonia uniflora, see Beadlily, queencup
Clobert, Jean, *see* Bro, Elisabeth
Clostridium botulinum, Type C, *see* Avian botulism
Clover: 575
Coccothraustes vespertinus, see Grosbeak, evening
Coccyzus americanus, see Cuckoo, yellow-billed
Cochran, Collins L. Jr., *see* Heffelfinger, James R.
Cocklebur: 359
Cocoplum: 874
Cocyzus americanus, see Cuckoo, yellow-billed
Coefficient of variation: 400, 800
Coffee, wild: 141
Colaptes auratus, see Flicker, northern
Colinus virginianus, see Quail, northern bobwhite
Collisions: 77, 81, 128
Collopy, Michael W., *see* Humes, Marcia L.
***Colobus satanus*:** 6–7
Colorado: 126–136, 270–277, 315–326, 470–483, 921–929, 1327–1336, 1345–1349
Colorado Desert: 159
Colubrina texensis, see Hogplum
Columba fasciata, see Pigeon, band-tailed
Columba leucocephala, see Pigeon, white-crowned
Columba livia, see Dove, rock
Columba livia, see Pigeon, feral
Columbia: 1–13
Columbina inca, see Dove, Inca
Columbina passerina, see Dove, common ground
Comptonia peregrina, see Fern, sweet
Condalia hookeri, see Brasil
Condor, Andean: 113
Condor, California: 112–113
Conecuh National Forest: 26–35
Confidence interval:
the insignificance of statistical significance testing, 763–772
Connecticut: 369–373
Conner, L. Michael, *see* Chamberlain, Michael J.
Conner, Mike, Bruce Plowman, Bruce D. Leopold, and Charles Lovell. Influence of time-in-residence on home range and habitat use of bobcats, 261–269
Conner, Richard N., *see* Carrie, N. Ross
Conocarpus erecta, see Buttonwood
Conover, Michael R., *see* Gosser, Allen L.
———, *see* Linnell, Michael A.
———, *see* Wagner, Kimberly K.
Conservation Reserve Program: 147–148, 189–198, 386, 420–440, 686–694, 705–710, 731, 977–989, 1299–1305
conservation reserve program: source or sink habitat for grassland birds in Missouri?, 530–538
effects of habitat on dickcissel abundance and nest success in conservation reserve program fields in Kansas, 523–529
Contopus borealis, see Flycatcher, olive-sided
Contopus sordidulus, see Peewee, western
Contopus virens, see Peewee, eastern
Conyza spp., *see* Mare's tail
Coot, American: 1233, 1363–1364
Copepods, cyclopoid: 362
Coragyps atratus, see Vulture, black
Cormorant, double-crested: 1016, 1336
Corn: 956–963, 980
Cornus florida, see Dogwood
Cornus nuttallii, see Dogwood, Pacific
***Cornus* spp:** 32
Cornus stolonifera, see Dogwood, red-osier
***Corophium* spp:** 17
Corvid: 445
Corvus brachrhynchos, see Raven, American
Corvus corax, see Raven, common
Corvus cryptoleucus, see Raven, Chihuahuan
Corylus cornuta, see Hazel, beaked
Corylus cornuta, see Hazel, California
Corynorhinus townsendii, see Bat, Townsend's big-eared
Costa Rica: 451
Coster, Dan, *see* Stolley, Dorie S.
Cottonwood: 78, 1120
Cottonwood, eastern: 630–636
Cottonwood, Fremont: 452–457
Cottonwood, plains: 459
Coturnix japonica, see Quail, Japanese
Cougar:
species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
winter prey selection by wolves and cougars in and near Glacier National Park, Montana, 901–910
Count, flush: 199–203
Count, point:
influence of point count length and repeated visits on habitat model performance, 815–823
***Couratari guianensis*:** 547
Cover, dense nesting:
effectiveness of dense nesting cover for increasing duck production in Saskatchewan, 382–389

Cowbird, brown-headed: 185–187, 489, 807, 1337
Cox, Robert R. Jr., *see* Lercel, Barbara A.
Coyote: 202, 270, 275, 278, 305, 386, 574–579, 615, 687, 731–738, 790, 890, 907, 936, 1104, 1300
effects of coyote removal on the faunal community in western Texas, 1066–1081
effect of preventive coyote hunting on sheep losses to coyote predation, 606–612
relative vulnerability of coyotes to removal methods on a northern California ranch, 939–949
species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
territoriality and breeding status of coyotes relative to sheep predation, 593–605
Coypu:
local determinants of coypu distribution along the Luján River, eastcentral Argentina, 895–900
Crakes, spotless: 122
Crane, sandhill: 1233
seasonal movements of sandhill cranes radiomarked in Yellowstone National Park and Jackson Hole, Wyoming, 126–136
Crane, whooping: 133
Crawford, John A., *see* Keegan, Thomas W.
Creatinine:
sampling and statistical analysis of snow-urine allantoin:cretinine ratios, 1118–1132
Creeper, brown: 187–188, 484–496
Creosote: 1182
***Cricetomys emini*:** 6–7
Cricket, Jerusalem: 931
Crinum americanum, see Lily, swamp
Crocodile, New Guinea: 1193
Crocodile, Nile: 1193
Crocodile, saltwater: 1193
Crocodylus niloticus, see Crocodile, Nile
Crocodylus novaeguineae, see Crocodile, New Guinea
Crocodylus porosus, see Crocodile, saltwater
Crocuta crocuta, see Hyena
Crop, cover: 426
Cropland: 675–685, 687
Crotalus adamanteus, see Rattlesnake, eastern diamondback
Croton: 1067
Croton spp., *see* Croton
Crowberry, low-lying: 335–336
CRP, *see* Conservation Reserve Program
Crypsis schoenoides, see Timothy, swamp
Cuckoo, yellow-billed: 188, 807, 816–821
Culicids: 362
Cupressus arizonica, see Cypress
Currant, sticky: 347
Cyanocitta cristata, see Jay, blue
Cyanocitta stelleri, see Jay, Steller's
Cygnus buccinator, see Swan, trumpter
Cygnus columbianus bewickii, see Swan, Bewick's
Cynomys leucurus, see Prairie dog, white-tailed
Cynomys ludovicianus, see Prairie dog, black-tailed
Cyperus erythrorhizis, see Sedge, redroot flat
Cypher, Brian L., and Nancy Frost. Condition of San Joaquin kit foxes in urban and exurban habitats, 930–938
Cypress: 452

D

Dabbert, C. Brad, *see* Mueller, James M.
Dactylis glomerata, see Grass, orchard
Dailey, Thomas V., *see* Frawley, Brian J.
Dalton, William J., *see* Lautenschlager, R. A.
Dama dama, see Deer, fallow
Dasyprocta spp., *see* Agouti
Dasypus novemcinctus, see Armadillo
David, Larry M., *see* Warner, Richard E.
Davis, J. Brian, *see* Blums, Peter
DDT: 791
Decker, David G., *see* Avery, Michael L.
Deer, Columbian black-tailed: 346–347, 600, 1333
movements, survival, and mortality of black-tailed deer in the Klickitat basin of Washington, 861–871
Deer, Chinese water: 1159
Deer, fallow: 567
Deer, grey brocket: 3
Deer, mule: 347, 862, 904, 923, 1327
mule deer survival in Colorado, Idaho, and Montana, 315–326
Deer, red: 342, 1090, 1110, 1128
real-time physiological responses of red deer to translocations, 1152–1162
Deer, red brocket: 3
Deer, roe: 1090, 1110
Deer, Japanese sika: 1090
Deer, white-tailed: 263, 278, 313, 346–347, 567, 613, 615, 901–910, 923, 935, 957, 965, 971, 1104, 1106, 1110, 1288, 1320–1326, 1332, 1336
effect of Hurricane Andrew on survival and movements of white-tailed deer in the everglades, 872–879
further evaluation of the genetic consequences of translocations on southeastern white-tailed deer population, 327–334
predation and survival of white-tailed deer fawns in northcentral New Brunswick, 574–579
residual effects of thinning and high white-tailed deer densities on northern redback salamanders in southern New england oak forests, 1172–1180
species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
survival and movements of orphaned white-tailed deer fawns in Texas, 570–574
Delany, David K., Teryl G. Grubb, Paul Beier, Larry L. Pater, and M. Hildegard Reiser. Effects of helicopter noise on Mexican spotted owls, 60–76
Delta Caribou Herd: 588–593
Demarais, Stephen, *see* Giuliano, William M.
———, *see* Mueller, James M.
DeMaynadier, Phillip G., and Malcolm L. Hunter, Jr.

Forest canopy closure and juvenile emigration by pool-breeding amphibians in Maine, 441–450
Demography: 1–13
demography of decline of the Red Wine Mountains caribou herd, 580–587
reintroduction and postrelease movements of red-cockaded woodpecker groups in eastern Texas, 824–832
survival of greater white-fronted geese: effects of year, season, sex, and body condition, 1239–1249
Denali National Park: 1105
Dendrocopos leucotos, see Woodpecker, European white-backed
Dendrocygna autumnalis, see Duck, black-bellied whistling
Dendroica caerulescens, see Warbler, black-throated blue
Dendroica coronata, see Warbler, yellow-rumped
Dendroica discolor, see Warbler, prairie
Dendroica dominica, see Warbler, yellow-throated
Dendroica fusca, see Warbler, blackburnian
Dendroica magnolia, see Warbler, magnolia
Dendroica nigrescens, see Warbler, black-throated gray
Dendroica occidentalis, see Warbler, hermit
Dendroica penslyvanica, see Warbler, chestnut-sided
Dendroica petechia, see Warbler, yellow
Dendroica pinus, see Warbler, pine
Dendroica striata, see Warbler, blackpoll
Dendroica virens, see Warbler, black-throated green
Denmark: 80
Density, brood:
abundance and productivity of forest songbirds in a managed, unfragmented landscape in Vermont, 180–188
Derksen, Dirk V., *see* Ward, David H.
Desmognathus fuscus, see Salamander, northern dusky
Dettmers, Randy, David A. Buehler, John G. Bartlett, and Nathan A. Klaus. Influence of point count length and repeated visits on habitat model performance, 815–823
Diadophis punctatus, see Snake, Key ringneck
Diadophis punctatus, see Snake, northern ringneck
Diceros bicornis michaeli, see Rhinoceros, black
Dickcissel: 530–538
effects of habitat on dickcissel abundance and nest success in conservation reserve program fields in Kansas, 523–529
Didelphis virginiana, see Opossum
Didymopanax morototoni, see Matchwood
Diervilla lonicera, see Honeysuckle
Diller, Lowell V., *see* Thome, Darrin M.
Dingo: 305
Diospyros texana, see Persimmon, Mexican
Dipodomys microps, see Rat, kanagroo
Dipodomys ordii, see Rat, Ord's kangaroo
Dipterans: 362
Distemper: 893, 930–938
Distichlis spicata, see Grass, spike
Disturbance:
effects of helicopter noise on Mexican spotted owls, 60–76
response of fall-staging brant and Canada geese to aircraft overflights in southwestern Alaska, 373–381
Dixie National Forest: 607
DNC, *see* Cover, dense nesting
Dog, domestic: 576, 687
Dog, wild: 906, 936
Dogwood: 309–314
Dogwood, Pacific: 292
Dogwood, red-osier: 1321
Dolbeer, Richard A., *see* Blackwell, Bradley F.
Dolichonyx oryzivorus, see Bobolink
Dove, common ground: 807
Dove, Inca: 81–81
Dove, mourning: 81–82, 232, 807, 1012–1013
molt of after-hatching-year mourning doves, 223–231
Dove, quail: 241
Dove, rock: 777
Dove, spotted: 999–1001
Dove, white-winged: 232
Dove, zebra: 999–1000
Dove, Zenaida:
population dynamics of Zenaida doves in Cidra, Puerto Rico, 232–244
Dredging:
use of dredged material substrates by nesting terns and black skimmers, 137–146
Drever, Mark C., *see* Hobson, Keith A.
Drew, John D., *see* Waithman, John D.
Drewien, Roderick C., Wendy M. Brown, John D. Varley, and David C. Lockman. Seasonal movements of sandhill cranes radiomarked in Yellowstone National Park and Jackson Hole, Wyoming, 126–136
Drift fence: 441–450
Dropseed: 1067
Drought:
effects of drought on desert tortoise movement and activity, 1181–1192
Dryocopus martius, see Woodpecker, black
Dryocopus pileatus, see Woodpecker, pileated
Duck, American black: 379, 393, 406, 647, 991, 1231
Duck, black-bellied whistling: 621, 1233
Duck, common shel: 621
Duck, Hawaiian: 999–1000
Duck, mallard: 382–389, 393, 418, 428–430, 489, 507–509, 630, 647, 657, 662, 1078, 1273, 1359–1369
environmental characteristics associated with the occurrence of avian botulism in wetlands of a northern California refuge, 358–368
mate loss in winter affects reproduction of mallards, 621–629

survival of postfledging mallards in northcentral Minnesota, 403–408
Duck, mottled: 991
Duck, mountain: 995
Duck, ruddy: 1362–1363
Duck, tufted: 657
Duck, wood: 404, 621
evaluation of a plasticine-filled leg band for day-old ducklings, 656–663
natural-cavity use by nesting wood ducks in Illinois, 630–638
wood duck brood movements and habitat use on prairie rivers in South Dakota, 639–649
Duda, Jeffery J., Anthony J. Krzysik, and Jerome E. Freilich. Effects of drought on desert tortoise movement and activity, 1181–1192
Duiker, blue: 5–7
Duiker spp.: 3, 5–7
Dumetella carolinensis, see Catbird, gray
Duncan, David C., *see* McKinnon, Donald T.

E

Eagle, bald: 71, 73, 95, 864, 958
double-survey estimates of bald eagle populations in Oregon, 794–802
movements and survival of bald eagles banded in the Greater Yellowstone Ecosystem, 781–793
Eagle, golden: 93
effects of fire on golden eagle territory occupancy and reproductive success, 773–780
Eagle, steppe: 111
Echinochloa crusgalli, see Grass, common barnyard
Echolocation: 553–561, 1308
Ecology, breeding:
abundance and productivity of forest songbirds in a managed, unfragmented landscape in Vermont, 180–188
comparative breeding ecology of Cooper's hawks in urban and exurban areas of southeastern Arizona, 77–84
site selection and nest success of ring-necked pheasants as a function of location in Iowa landscapes, 976–989
Ecology, foraging:
caribou calf production and seasonal range quality during a population decline, 335–345
characteristics of foraging trees selected by pileated woodpeckers in New Brunswick, 361–469
conifer release alternatives increase aspen forage quality in northwestern Ontario, 1320–1326
effects of summer sheep grazing on browse nutritive quality in autumn and winter, 346–354
estimation of habitat selection for central-place foraging animals, 1028–1038
resource variability and conservation of American flamingos in coastal wetlands of Yucatán, Mexico, 1201–1212
selection of pines for foraging by red-cockaded woodpeckers, 843–852
Ecology, landscape: 431–440
abundance and productivity of forest songbirds in a managed, unfragmented landscape in Vermont, 180–188
slack in the configuration of habitat patches for northern bobwhites, 245–250
survival of ring-necked pheasant hens during spring in relation to landscape features, 147–154
Ecology, riparian: 167–179
Ecuador: 1–13
Eddleman, William R., *see* Legare, Michael L.
Edge: 147–154, 523–529, 980
Egg peritonitis: 506
Eider, common: 662
El Salvador: 451
Elaphe obsoleta obsoleta, see Snake, black rat
Elaphe obsoleta spiloides, see Snake, gray rat
Elderberry: 485
Electrical conductivity:
use of total body electrical conductivity to predict northern bobwhite lipid mass, 695–704
Electricity:
effects of deterrents on avian abundance and nesting density in electrical substations in Oklahoma, 1009–1017
Eleocharis spp., *see* Rush, spike
Elephant, African: 794
Elk: 787, 901–910, 923, 950, 1101, 1132
adaptive fetal sex allocation in elk: evidence and implications, 1109–1117
effectiveness of Hot Sauce® and Deer Away® repellents for deterring elk browsing of aspen sprouts, 1327–1336
species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
sampling and statistical analysis of snow-urine allantoin:cretinine ratios, 1118–1132
Elk, Rocky Mountain: 346–347, 567–568, 594
Elk, Roosevelt: 347, 1109–1117
Elk, tule:
use of fecal steroid metabolites to estimate the pregnancy rate of a free-ranging herd of tule elk, 561–569
Ellsworth, Darrell L., *see* Leberg, Paul L.
Elm, American: 640
Ely, Craig R., *see* Schmutz, Joel A.
Embrotoxicity:
selenium accumulation and effects on reproduction in captive American kestrels fed selenomethionine, 502–511
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
Emigration: 580–587
forest canopy closure and juvenile emigration by pool-breeding amphibians in Maine, 441–450
Empetrum nigrum, see Crowberry, low-lying
Empidonax difficilis, see Flycatcher, Pacific-slope

Empidonax flaviventris, see Flycatcher, yellow-bellied
Empidonax fulvifrons, see Flycatcher, buff-breasted
Empidonax hammondii, see Flycatcher, Hammond's
Empidonax traillii, see Flycatcher, willow
Empidonax virescens, see Flycatcher, Acadian
Emydoidea blandingii, see Turtle, Blanding's
Endangered Species Act: 470–483, 511, 921, 1181
Energetics:
energy-based carrying capacity for quails, 664–674
population energetics of northern pintails wintering in the Sacramento Valley, California, 1222–1238
England: 374, 708–709, 933
Enhydra lutris, see Otter, sea
Enterobacter spp.: 207
Enzyme immunoassay:
use of fecal steroid metabolites to estimate the pregnancy rate of a free-ranging herd of tule elk, 561–569
***Eperua falcata*:** 547
Ephydra spp., *see* Fly, brine
Ephydrids: 362
Epidonax minimus, see Flycatcher, least
Epilobium spp., *see* Fireweed
Eptesicus fuscus fuscus, see Bat, eastern big brown
Eptesicus fuscus, see Bat, big brown
Equisetum arvense, see Horsetail, field
Equus burchelli, see Zebra
Eragrostis maicense, see Grass, love
Eremophila alpestris, see Lark, horned
Erickson, Wallace P., *see* Ward, David H.
Ericsson, Goran, *see* Swenson, Jon E.
Eriophorum angustifolium, see Grass, cotton
Erythrocebus patas, see Monkey, patas
Esker, Terry L., *see* Herkert, James R.
Etter, Stanley L., *see* Warner, Richard E.
Eubalaena glacialis, see Whale, right
Eubranchipods: 362
Eudocimus albus, see Ibis, white
Euliss, Ned H. Jr., *see* Rocke, Tonie E.
Euphagus carolinus, see Blackbird, rusty
Eurycea bislineata, see Salamander, northern two-lined
Evaporation basin:
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
Everglades National Park: 872–879
Everlasting: 347
Ewalt, Darl R., *see* Roffe, Thomas J.
Exotic species:
effective monitoring for adaptive wildlife management: lessons from the Galápagos Islands, 1055–1065
correlated factors in amphibian decline: exotic species and habitat change in western Washington, 1162–1171
northern bobwhite chick mortality caused by red imported fire ants, 1291–1298
norway rats as predators of burrow-nesting seabirds: insights from stable isotope analyses, 14–25
range expansion, population sizes, and management of wild pigs in California, 298–308
Extripation: 1–13, 333
Eysenhardtia texana, see Kidneywood, Texas

F

Fagus grandifolia, see Beech, American
Fagus sylvatica, see Beech
Falco columarius, see Merlin
Falco mexicanus, see Falcon, prairie
Falco peregrinus, see Falcon, peregrine
Falco rusticolus, see Gyrfalcon
Falco sparvarius, see Kestrel, American
Falcon, peregrine: 81
performance and accuracy evaluation of small satellite transmitters, 1349–1358
Falcon, prairie: 72–73, 1067
Fecal analysis: 922, 561–569
Fecal steroids:
use of fecal steroid metabolites to estimate the pregnancy rate of a free-ranging herd of tule elk, 561–569
Feeding, supplemental:
supplemental feeding regimes for Egyptian vultures in the Negev Desert, Israel, 107–115
Feline panleukopenia: 614–619
Felis concolor coryi, see Panther, Florida
Fern, bracken: 347, 443, 575
Fern, sweet: 443
Fern, sword: 293, 486
Ferret, black-footed: 271
Fescue, Idaho: 1119–1120
Fescue, meadow: 518
Fescue, sheep: 347
Festuca idahoensis, see Fescue, Idaho
Festuca ovina, see Fescue, sheep
Festuca pratensis, see Fescue, meadow
Ficus aurea, see Fig, strangler
Fig, strangler: 874
Fimbristylis castanea, see Sedge, tropical
Finch, bull: 227
Finch, house: 305, 999–1000, 1013
Finch, purple: 188, 489–490
Fir, balsam: 167–179, 181, 461–468, 575, 581, 1307, 1321
Fir, big-cone: 45
Fir, Douglas: 36–37, 45, 57, 61, 99, 205, 292, 317, 347, 452, 485, 492, 553–554, 795, 863, 902, 1083, 1120, 1213, 1315, 1377, 1382
Fir, grand: 99, 347, 795, 863
Fir, Pacific silver: 863
Fir, subalpine: 316–317, 863, 902, 1083, 1328
Fir, white: 36–37, 45
Fire: 459, 519, 527, 535, 538, 693, 803, 859, 1382
effects of fire on golden eagle territory occupancy and reproductive success, 773–780

Firebush: 309–314
Fireweed: 347
Fischer: 576
Fish Springs National Wildlife Refuge: 990–996
Fishlake National Forest: 607
Flag: 874
Flake, Lester D., *see* Gabbert, Andy E.
———, *see* Granfors, Diane A.
Flamingo, American:
resource variability and conservation of American flamingos in coastal wetlands of Yucatán, Mexico, 1201–1212
Flamingo, Chilean: 1209
Flathead National Forest: 902, 1083
Flemming, Stephen P., Gillian L. Holloway, E. Jane Watts, and Peter S. Lawrance. Characteristics of foraging trees selected by pileated woodpeckers in New Brunswick, 461–469
Flicker, northern: 170–177, 188, 489–490
Florida: 33, 116–125, 251–260, 328, 619, 782, 833–842, 872–879, 1004, 1193–1200
Florida Everglades:
effect of Hurricane Andrew on survival and movements of white-tailed deer in the everglades, 872–879
Fly, brine: 1209
Fly, nasal bot: 588–593
Fly, warble: 588–593
Flyatcher, Acadian: 807
Flycatcher, buff-breasted: 459
Flycatcher, great-crested: 184–188, 807
Flycatcher, Hammond's: 484–496
Flycatcher, least: 184–188
Flycatcher, olive-sided: 170–177, 188, 490
Flycatcher, Pacific-slope: 484–496
Flycatcher, scissor-tailed: 1013
Flycatcher, willow: 457, 489–490
Flycatcher, yellow-bellied: 170–177
Flyway, Atlantic: 652, 662
Flyway, Central: 639–649, 652
Flyway, Mississippi: 393, 621–629, 630, 652, 662
Flyway, Pacific: 1239–1249
Food web: 14–25
Forbes, Andrew R., *see* Mueller, James M.
Forbes, Graham J., *see* Ballard, Warren B.
Forest management:
abundance and productivity of forest songbirds in a managed, unfragmented landscape in Vermont, 180–188
bat activity in thinned, unthinned, and old-growth forests in western Oregon, 553–561
breeding bird assemblages inhabiting riparian buffer strips in Newfoundland, Canada, 167–179
forest canopy closure and juvenile emigration by pool-breeding amphibians in Maine, 441–450
forest stand characteristics and reproduction of northern spotted owls in managed north-coastal California forests, 44–59
habitat selection by forest bats in relation to mixed-wood stand types and structure in central Ontario, 1306–1319
influence of riparian buffer width on bird assemblages in western Oregon, 484–496
movements of northern flying squirrels in different-aged forest stands of western Oregon, 291–297
relations of small mammal populations to even-aged shelterwood systems: a comment, 1376–1380
relations of small mammal populations to even-aged shelterwood systems: a reply, 1381–1389
residual effects of thinning and high white-tailed deer densities on northern redback salamanders in southern New england oak forests, 1172–1180
scrub-successional bird community dynamics in young and mature longleaf pine-wiregrass savannahs, 803–814
Forest, old-growth: 39, 291–297, 486, 824, 843–852, 1306–1319
bat activity in thinned, unthinned, and old-growth forests in western Oregon, 553–561
old-forest distribution around spotted owl nests in the central Cascade Mountains, Oregon, 1212–1221
Fort Niobrara National Wildlife Refuge: 950–955
Forys, Elizabeth A., and Stephen R. Humphrey. Use of population viability analysis to evaluate management options for the endangered Lower Keys marsh rabbit, 251–260
Fouquieria splendens, see Ocotillo
Fowl, mallee: 1147
Fox, gray: 615, 933, 1066–1081
Fox, red: 111, 147, 202, 405, 575, 719, 731–738, 933, 985, 987, 1046, 1077, 1300
Fox, San Joaquin kit:
condition of San Joaquin kit foxes in urban and exurban habitats, 930–938
Fox, swift: 270, 275
Frampton, Chris M., *see* Henderson, Ray J.
France: 511–516, 1044–1051
Frawley, Brian J., David A. Osborne, Harmon P. Weeks, Loren W. Burger, and Thomas V. Dailey. Use of total body electrical conductivity to predict northern bobwhite lipid mass, 695–704
Fraxinus nigra, see Ash, black
Fraxinus pennsylvanica, see Ash, green
Fraxinus velutina, see Ash, velvet
Freilich, Jerome E., *see* Duda, Jeffery J.
French Guiana: 546–552
Fritzell, Erik K., *see* Gehrt, Stanley D.
Frog, bull:
correlated factors in amphibian decline: exotic species and habitat change in western Washington, 1162–1171
Frog, Pacific tree: 1169
Frog, pickerel: 1175
Frog, red-legged:
correlated factors in amphibian decline: exotic species and habitat change in western Washington, 1162–1171

Frog, wood: 441–450, 1172
Frost, Nancy, *see* Cypher, Brian L.
Fulica americana, see Coot, American
Functional response:
power analysis of wolf-moose functional responses, 396–402
***Funisciurus* spp:** 6–7

G

Gabbert, Andy E., Anthony P. Leif, Jon R. Purvis, and Lester D. Flake. Survival and habitat use by ring-necked pheasants during two disparate winters in South Dakota, 711–722
Gaddis, Stanley E., *see* Shumake, Stephen A.
Gadwall: 382–389, 647, 657, 1359–1369
Galápagos Islands:
effective monitoring for adaptive wildlife management: lessons from the Galápagos Islands, 1055–1065
Gallinula chloropus, see Moorhen, common
Gambelia sila, see Lizard, blunt-nosed leopard
Gardner, Ian A., *see* Waithman, John D.
Garner, Dale L., *see* Hubbard, Michael W.
Garrett, Monte G, *see* Anthony, Robert G.
Garrot, Robert A., *see* Pils, Andrew C.
Garrya wrightii, see Silktassel
Gastropods: 362
Gau, Robert J., and Ray Case. Evaluating nutritional condition of grizzly bears via select blood parameters, 286–291
Gaultheria shallon, see Salal
Gava immer, see Loon, common
Gazella thomsoni, see Gazella
Gazelle: 906
Gehrt, Stanley D., and Erik K. Fritzell. Survivorship of a nonharvested raccoon population in south Texas, 889–894
Genetic drift: 253
Geochelone nigra, see Tortoise, Galápagos
Geographic Information System: 46–48, 148, 180–181, 189–198, 212–222, 254–260, 263, 298–308, 422, 631, 713, 733, 774, 845, 979–989, 1063, 1190, 1214, 1351
a geographic information system method for estimation home range size, 748–755
Geopelia striata, see Dove, zebra
Georgia: 32, 538–545, 805
Geothlypis trichas, see Yellowthroat, common
Geotrygon spp., *see* Dove, quail
Geranium richardsoni, see Geranium, white
Geranium, white: 1328
Giajillo: 570
Gibbs, James P., Howard L. Snell, and Charlotte E. Causton. effective monitoring for adaptive wildlife management: lessons from the Galápagos Islands, 1055–1065—Invited paper
Gibeau, Michael L., *see* McLellan, Bruce N.
Gidlewski, Thomas, *see* Roffe, Thomas J.
Gila National Forest: 38
GIS, *see* Geographic Information System
Gitzen, Robert A., *see* Seaman, D. Erran
Giuliano, William M., Stephen Demarais, Robert E. Zaiglin, and Misty L. Sumner. Survival and movements of orphaned white-tail deer fawns in Texas, 570–574
Glacier National Park: 901–910, 921–929, 1082–1093, 1094–1108
Glaucomys sabrinus, see Squirrel, northern flying
Glaucomys volans, see Squirrel, southern flying
Global Positioning System: 65, 88, 117, 272, 816, 864, 1042, 1063, 1183, 1214
Gnatcatcher, blue-gray: 807, 812
Goldeneye, Barrow's: 1359–1369
Goldenrod: 141, 855
Goldfinch, American: 187, 488–491, 530–538, 807
Goose, barnacle: 379, 621, 654
Goose, cackling Canada: 1244
Goose, Canada: 227, 422–440, 621, 1363–1364
effects of saline drinking water on early gosling development, 990–996
molt migrant Canada geese in northern Ontario and western James Bay, 649–655
plant growth regulator (Stronghold®) enhances repellency of anthraquinone formulation (Flight Control®) to Canada geese, 1336–1343
response of fall-staging brant and Canada geese to aircraft overflights in southwestern Alaska, 373–381
will the availability of insular nesting sites limit reproduction in urban Canada goose populations?, 369–373
Goose, emperor: 1244
Goose, giant Canada: 370
molt migrant Canada geese in northern Ontario and western James Bay, 649–655
Goose, greater snow: 374, 1244, 1287
Goose, greater white-fronted:
survival of greater white-fronted geese: effects of year, season, sex, and body condition, 1239–1249
Goose, Greenland white-fronted: 1246
Goose, lesser snow: 379, 621, 653
Goose, pink-footed: 379, 654
Goose, Ross's: 621
Goose, western Canada: 991
molt migrant Canada geese in northern Ontario and western James Bay, 649–655
Gopher, northern pocket:
repellents to reduce cable gnawing by northern pocket gophers, 1344–1349
Gopherus agassizii, see Tortoise, desert
Gordus, Andrew G. Selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
Goshawk: 728
dawn and morning goshawk courtship vocalizations as a method for detecting nest sites, 511–516

efficacy of northern goshawk broadcast surveys in Washington state, 98–106
Gosser, Allen L., and Michael R. Conover. Will the availability of insular nesting sites limit reproduction in urban Canada goose populations?, 369–373
GPS, *see* Global Positioning System
Grackle, boat-tailed:
seed color avoidance by captive red-winged blackbirds and boat-tailed grackles, 1003–1008
Grackle, common: 1011, 1014
Graham, Jeri L., *see* Lautenschlager, R. A.
Grama, black: 1067
Grama, blue: 86, 1067, 1300
Grama, sideoats: 1300
Grand Teton National Park: 130, 782, 787
Granfors, Diane A., and Lester D. Flake. Wood duck brood movements and habitat use on prairie rivers in South Dakota, 639–649
Granjeno: 571
Grape, Oregon: 293, 486, 555
Grass, big bluestem: 518, 524, 687, 1300
Grass, blue: 347, 518, 1328
Grass, blue-bunch wheat: 316–317, 1119–1120
Grass, bluestem: 26–27, 29, 855
Grass, buffalo: 1300
Grass, cheat: 271, 316, 773
Grass, common barnyard: 1261–1272
Grass, cord: 117, 251, 254
Grass, cotton: 336
Grass, eel: 374
Grass, green bristle: 1269–1272
Grass, Guinea: 233
Grass, hooded windmill: 1067
Grass, Indian: 518, 524, 687, 1300
Grass, Indian rice: 271
Grass, little bluestem: 518, 524, 687, 1300
Grass, love: 874
Grass, muhly: 874
Grass, musk: 1203
Grass, orchard: 518
Grass, panic: 141, 1261–1272
Grass, plains bluestem: 1300
Grass, redtop bent: 518
Grass, reed: 16–17
Grass, sand cord: 874
Grass, Sandberg blue: 271
Grass, saw: 117, 873
Grass, spike: 117
Grass, switch: 518, 524, 687, 712, 980
Grass, western wheat: 86, 524
Grass, wheat: 272
Grass, widgeon: 1203
Grass, wire: 26–27, 29, 34
scrub-successional bird community dynamics in young and mature longleaf pine-wiregrass savannahs, 803–814
Grasshopper: 931
Grassland: 429, 639–649, 675–685, 731, 895–900, 958–963, 982–984, 1066–1081, 1273–1281
conservation reserve program: source or sink habitat for grassland birds in Missouri?, 530–538
effects of habitat on dickcissel abundance and nest success in conservation reserve program fields in Kansas, 523–529
response of northern harriers and short-eared owls to grassland management in Illinois, 517–523
Gray, Matthew J., Richard M. Kaminski, and Govinda Weerakkody. Predicting seed yield of moist-soil plants, 1261–1268
———, and Michael G. Brasher. A new method to predict seed yield of moist-soil plants, 1269–1272
Grazing: 346–354, 423, 535, 693, 859, 1120, 1336–1343
Greasewood, black: 86, 271, 778
Greater Yellowstone Ecosystem:
movements and survival of bald eagles banded in the Greater Yellowstone Ecosystem, 781–793
trophic relations of brown and black bears in several North American ecosystems, 921–929
Greater Yellowstone Interagency Brucellosis Committee: 950, 1133
Green Mountain National Forest: 180–188
Greenbrier: 877
Griffon, Cape: 112
Griffon, Eurasian: 107–108, 111–112
Grosbeak, black-headed: 489–490
Grosbeak, blue: 806–809
Grosbeak, evening: 187, 188, 489
Grosbeak, pine: 170–177, 188
Grosbeak, rose-breasted: 184–188
Grouse, black: 723
Grouse, red: 163
Grouse, ruffed: 188, 723, 1287
Grouse, sage: 271
Grouse, willow:
mortality causes and survival rates of hunted and unhunted willow grouse, 722–730
Grubb, Teryl G., *see* Delany, David K.
Grus americana, see Crane, whooping
Grus canadensis tabida, see Crane, sandhill
Gryllacrididae, see Cricket, Jerusalem
Guam: 1015
Guatemala: 451
Guichón, María Laura, and Marcelo Hernán Cassini. Local determinants of coypu distribution along the Luján River, eastcentral Argentina, 895–900
Guiraca caerulea, see Grosbeak, blue
Gull, glaucous-winged: 990
Gulo gulo, see Wolverine
Gum, black: 1282
Gum, sweet: 1282
***Gustavia augusta*:** 547
Guthery, Fred S. Energy-based carrying capacity for quails, 664–674
——— Slack in the configuration of habitat patches for Northern Bobwhites, 245–250

———, *see* Heffelfinger, James R.
Gutiérrez, R. J., *see* Peery, M. Zachariah
Guyer, Craig, *see* Aresco, Matthew J.
GYIBC, *see* Greater Yellowstone Interagency Brucellosis Committee
Gymnogyps californianus, see Condor, California
Gypaetus barbatus, see Vulture, bearded
Gyps coprotheres, see Griffon, Cape
Gyps fulvus, see Griffon, Eurasian
Gyrfalcon: 71, 728

H

Haas, Carola A, *see* Carter, Shawn L.
Habitat fragmentation: 276, 441–450, 461, 1212–1221
Habitat, nesting:
effectiveness of dense nesting cover for increasing duck production in Saskatchewan, 382–389
natural-cavity use by nesting wood ducks in Illinois, 630–638
use of dredged material substrates by nesting terns and black skimmers, 137–146
will the availability of insular nesting sites limit reproduction in urban Canada goose populations?, 369–373
Habitat selection: 461–469
conservation reserve program: source or sink habitat for grassland birds in Missouri?, 530–538
estimation of habitat selection for central-place foraging animals, 1028–1038
habitat composition and configuration around Mexican spotted owl nest and roost sites in the Tularosa Mountains, New Mexico, 36–43
habitat selection by forest bats in relation to mixed-wood stand types and structure in central Ontario, 1306–1319
habitat use of eastern wild turkeys in central Mississippi, 210–222
home range and habitat selection of bog turtles in southwestern Virginia, 853–860
influence of time-in-residence on home range and habitat use of bobcats, macrohabitat effects on summer survival, movements, and clutch success of northern bobwhite in Kansas, 675–685
microhabitat selection by nesting and brood-rearing northern bobwhite in Kansas, 686–694
multiscaled habitat selection by elegant trogons in southeastern Arizona, 451–461
old-forest distribution around spotted owl nests in the central Cascade Mountains, Oregon, 1212–1221
response of northern harriers and short-eared owls to grassland management in Illinois, 517–523
site selection and nest success of ring-necked pheasants as a funnction of location in Iowa landscapes, 976–989
slack in the configuration of habitat patches for northern bobwhites, 245–250
survival and habitat use by ring-necked pheasants during two diparate winters in South Dakota, 711–722
survival of ring-necked pheasant hens during spring in relation to landscape features, 147–154
wood duck brood movements and habitat use on prairie rivers in South Dakota, 639–649
Habitat, urban:
will the availability of insular nesting sites limit reproduction in urban Canada goose populations?, 369–373
Hagar, Joan C. Influence of riparian buffer width on bird assemblages in western Oregon, 484–496
Haliaeetus leucocephalus, see Eagle, bald
Hall, Linnea S., and R. William Mannan. Multiscaled habitat selection by elegant trogons in southeastern Arizona, 451–461
Hanley, Thomas A., *see* Jacoby, Michael E.
Hansen, Leslie A., *see* Lee, Bruce A. Vander
***Haplotrema* spp:** 17
Harmata, Alan R., George J. Montopoli, Bob Oakleaf, Peter J. Harmata, and Marco Restani. Movements and survival of bald eagles banded in the Greater Yellowstone Ecosystem, 781–793
Harmata, Peter, J., *see* Harmata, Alan R.
***Harpaphe* spp: 17**
Harrier, northern: 507, 1067, 1273–1281
response of northern harriers and short-eared owls to grassland management in Illinois, 517–523
Harrington, Fred H., *see* Schaefer, James A.
Hartless, Christine S., *see* Labisky, Ronald F.
Harvest: 163–165, 570–574, 861–871, 880–888, 895–900, 956–963, 1087–1088, 1111, 1116, 1123
effect of preventive coyote hunting on sheep losses to coyote predation, 606–612
effects of coyote removal on the faunal community in western Texas, 1066–1081
effects of egg and hatchling harvest on American alligators in Florida, 1193–1200
effects of fall hunting on wild turkey populations in Virginia and West Virginia, 964–975
mate loss in winter affects reproduction of mallards, 621–629
mortality causes and survival rates of hunted and unhunted willow grouse, 722–730
survivorship of a nonharvested raccoon population in south Texas, 889–894
towards wildlife management in tropical forests, 1–13
Harvest, timber: 167–179
Havera, Stephen P., *see* Yetter, Aaron P.
Hawaii: 997–1003, 1056
Hawk, broad-winged: 1067
Hawk, Cooper's: 202, 513
comparative breeding ecology of Cooper's hawks in urban and exurban areas of southeastern Arizona, 77–84
Hawk, ferruginous: 1067

an aerial sightability model for estimating ferruginous hawk population size, 85–97
Hawk, Harris': 1067
Hawk, red-tailed: 188, 202, 489–490, 507, 1067, 1273–1281
Hawk, sharp-shinned: 1067
Hawk, sparrow: 80, 778
Hawk, Swainson's: 1067
Hayes, John P., *see* Humes, Marcia L.
Haying: 519
Hays, David W., *see* Watson, James W.
Hazel, beaked: 1307, 1320–1326
Hazel, California: 293
Heffelfinger, James R., Fred S. Guthery, Ronald J. Olding, Collins L. Cochran, Jr., and Craig M. McMullen. Influence of precipitation timing and summer temperatures on reproduction of Gambel's quail, 154–161
Helicopter:
effects of helicopter noise on Mexican spotted owls, 60–76
response of fall-staging brant and Canada geese to aircraft overflights in southwestern Alaska, 373–381
***Heliosciurus* spp:** 6–7
Hellgren, Eric C., *see* James, John B.
Helmitheros vermivorous, see Warbler, worm-eating
Hemidactylium scutatum, see Salamander, four-toed
***Hemigrapsus* spp:** 17
Hemlock, eastern: 181, 442, 575, 816, 1174
Hemlock, mountain: 863
Hemlock, western: 16, 99, 292, 485, 554, 863, 1213
Henderson, Ray J., Chris M. Frampton, David R. Morgan, and Graham J. Hickling. The efficacy of baits containing 1080 for control of brushtail possums, 1138–1151
Henke, Scott E., and Fred C. Bryant. Effects of coyote removal on the faunal community in western Texas, 1066–1081
Hennager, Steven G., *see* Roffe, Thomas J.
Henry, Olivier, *see* Judas, Jacky
Hepatitus, infectious canine: 920–938
Herkert, James R., Scott A. Simpson, Ronald L. Westemeier, Terry L. Esker, and Jeff W. Walk. Response of northern harriers and short-eared owls to grassland management in Illinois, 517–523
Heron, black-crowned night: 508, 1016, 1274
Heron, great blue: 1016
Heske, Edward J., *see* Mosillo, Maia
Heterogeneity:
direct recovery rates of lesser scaup banded in northwest Minnesota: sources of heterogeneity, 389–395
Hibernation: 1189
Hickling, Graham J., *see* Henderson, Ray J.
Hickory, mockernut: 1282
Hickory, shagbark: 199, 731, 965
Hilderbrand, Grant V., *see* Jacoby, Michael E.
Himantopus mexicanus, see Stilt, black-necked
Hine, Christopher S., *see* Yetter, Aaron P.
Hobson, Keith A., Mark C. Drever, and Gary W. Kaiser. Norway rats as predators of burrow-nesting seabirds: insights from stable isotope analyses, 14–25
Hodges, Kurt M., *see* Chamberlain, Michael J.
Hogplum: 571
Hollick, John, *see* Mason, J. Russell
Holloway, Gillian L., *see* Flemming, Stephen P.
Holly, dahoon: 874, 877
Holodiscus discolor, see Oceanspray
***Homalomena pictuata*:** 547
Home range: 151, 428, 456, 639–649, 675–685, 731–738, 880–888, 1097–1098, 1181–1192
a geographic information system method for estimation home range size, 748–755
autocorrelation of location estimates and the analysis of radiotraking data, 1039–1044
effect of Hurricane Andrew on survival and movements of white-tailed deer in the everglades, 872–879
effects of fire on golden eagle territory occupancy and reproductive success, 773–780
effects of sample size on kernel home range estimates, 739–747
home range and habitat selection of bog turtles in southwestern Virginia, 853–860
influence of time-in-residence on home range and habitat use of bobcats, 261–269
movements of northern flying squirrels in different-aged forest stands of western Oregon, 291–297
seasonal variation of home range of collared peccary in tropical rain forests of French Guiana, 546–552
site selection and nest success of ring-necked pheasants as a function of location in Iowa landscapes, 976–989
territoriality and breeding status of coyotes relative to sheep predation, 593–605
Honduras: 451
Honeysuckle: 310
Hornocker, Maurice G., *see* Kunkel, Kryan E.
Horsetail, field: 336, 912
Hovey, Fred W., *see* McLellan, Bruce N.
Hubbard, Michael W., Dale L. Garner, Erwin E. Klaas. Wild turkey poult survival in southcentral Iowa, 199–203
———, Dale L. Garner, and Erwin E. Klaas. Factors influencing wild turkey hen survival in southcentral Iowa, 731–738
Huber, Royce, *see* Roffe, Thomas J.
Huckleberry: 293, 317, 486, 912, 1120
Hughes, John P., Robert J. Robel, Kenneth E. Kemp, and John L. Zimmerman. Effects of habitat on dickcissel abundance and nest success in conservation reserve program fields in Kansas, 523–529
Huisache: 571, 1292
Humes, Marcia L., John P. Hayes, and Michael W.

Collopy. Bay activity in thinned, unthinned, and old-growth forests in western Oregon, 553–561
Hummingbird, black-chinned: 227
Hummingbird, ruby-throated: 188, 227, 807
Hummingbird, rufous: 484–496
Humphrey, John S., *see* Avery, Michael L.
Humphrey, Stephen R., *see* Forys, Elizabeth A.
Hunter, Malcolm L., Jr., *see* DeMaynadier, Phillip G.
Hunting: 163–165, 389–395, 403–408, 570–574, 861–871, 880–888, 911–920, 956–963, 1087–1088, 1111, 1116, 1123, 1240
effect of preventive coyote hunting on sheep losses to coyote predation, 606–612
effects of coyote removal on the faunal community in western Texas, 1066–1081
effects of fall hunting on wild turkey populations in Virginia and West Virginia, 964–975
mate loss in winter affects reproduction of mallards, 621–629
mortality causes and survival rates of hunted and unhunted willow grouse, 722–730
survivorship of a nonharvested raccoon population in south Texas, 889–894
towards wildlife management in tropical forests, 1–13
Hunting, subsistence: 1–13, 861–871
Hurricane: 1201–1212
effect of Hurricane Andrew on survival and movements of white-tailed deer in the everglades, 872–879
Hurst, George A., *see* Miller, Darren A.
Hyaena hyaena, see Hyena, striped
Hydrilla: 1196
Hydropotes inermis, see Deer, Chinese water
Hyemoschus aquaticus, see Duiker spp.
Hyena: 907
Hyena, striped: 109, 111
Hyla crucifer crucifer, see Peeper, northern spring
Hyla regilla, see Frog, Pacific tree
Hylocichla mustelina, see Thrush, wood
Hypoderma tarandi, see Fly, warble
Hypothesis, condition-bias: 389–395

I

Ibis, white: 1210
Ictalurus spp., *see* Bullhead
Icteria virens, see Chat, Yellow-breasted
Icterus spurius, see Oriole, orchard
Ictinia mississippiensis, see Kite, Mississippi
Idaho: 104, 126–136, 315–326, 346–354, 606–612, 773–780, 782–790, 901, 911–929, 950–955, 1103, 1106
Ilex cassine, see Holly, dahoon
***Ilex* spp:** 29, 32
Illinois: 278–286, 470–483, 517–523, 630–638, 695–710, 956–963, 986
Implantation, intramuscular: 165
Implantation, subcutaneous: 162–166
Incubation: 33
Indian poke: 575
Indiana: 695–704
Indonesia: 6, 1193
Ingram, John R., *see* Waas, Joseph R.
Insects: 399–400
Interspersion: 249
Invertabrates: 358–368, 362, 708, 1205–1210, 1225, 1250
Iowa: 147–154, 199–203, 470–483, 534, 731–738, 887, 889–894, 972, 976–989
Isaacs, Frank B., *see* Anthony, Robert G.
Islands:
use of dredged material substrates by nesting terns and black skimmers, 137–146
will the availability of insular nesting sites limit reproduction in urban Canada goose populations?, 369–373
Isotope analyses: 267, 921–929
norway rats as predators of burrow-nesting seabirds: insights from stable isotope analyses, 14–25
Israel: 107–115
Italy: 511
Ixoreus naevius, see Thrush, varied

J

Jackal, black-backed: 945
Jackson Hole:
seasonal movements of sandhill cranes radiomarked in Yellowstone National Park and Jackson Hole, Wyoming, 126–136
Jacoby, Michael E., Grant V. Hilderbrand, Christopher Servheen, Charles C. Schwartz, Stephen M. Arthur, Thomas A. Hanley, Charles T. Robbins, and Robert Michener. Trophic relations of brown and black bears in several North American ecosystems, 921–929
Jaeger, Michael M., *see* Sacks, Benjamin N.
Jaguar: 549, 907
James, John B., Eric C. Hellgren, and Ronald E. Masters. Effects of deterrents on avian abundance and nesting density in electrical substations in Oklahoma, 1009–1017
Jasper National Park: 912, 1094
Jay, blue: 187–188, 807
Jay, Florida scrub:
evaluation of translocation for the threatened Florida scrub-jay, 833–842
Jay, gray: 105, 170–177
Jay, Steller's: 105, 488–492
Jenkins, Roger A., *see* Ballard, Warren B.
Jennings, Michael L., *see* Rice, Kenneth G.
Jensen, Allen E., *see* Roffe, Thomas J.
***Jessenia bataua*:** 547
John F. Kennedy International Airport: 1001
Johnson, Douglas H. The insignificance of statistical significance testing, 763–772—Invited paper

Johnson, James E., *see* Thogmartin, Wayne E.
Johnston, John J., *see* Mason, J. Russell
Judas, Jacky, and Olivier Henry. Seasonal variation of home range of collared peccary in topical rain forests of French Guiana, 546–552
Juglans major, see Walnut, black
Junco, dark-eyed: 170–177, 187–188, 490
Junco hyemalis, see Junco, dark-eyed
Juncus roemerianus, see Needlerush
Jung, Thomas S., Ian D. Thompson, Rodger D. Titman, and Andrew P. Applejohn. Habitat selection by forest bats in relation to mixed-wood stand types and structure in central Ontario, 1306–1319
Juniper: 86, 795
Juniper, alligator: 37, 452
Juniper, Rocky Mountain: 317
Juniper, Utah: 86, 316
Juniperus deppeana, see Juniper, alligator
Juniperus osteosperma, see Juniper, Utah
Juniperus scopulorum, see Juniper, Rocky Mountain

K

Kadlec, John A., *see* Stolley, Dorie S.
Kaiser, Gary W., *see* Hobson, Keith A.
Kaminski, Richard M., *see* Gray, Matthew J.
———, *see* Lercel, Barbara A.
Kansas: 470–483, 523–529, 535, 675–694, 735
Kaplan-Maier method:
effects of violation of assumptions for survival analysis methods in radiotelemetry studies, 1369–1375
Kasworm, Wayne F., *see* McLellan, Bruce N.
Keegan, Thomas W., and John A. Crawford. Reproduction and survival of Rio Grande turkeys in Oregon, 204–210
Kelly, Colleen, *see* Legare, Michael L.
Kemp, Kenneth E., *see* Hughes, John P.
Kenai Peninsula: 921–929
Kennedy, Patricia L., *see* Britten, Michael W.
Kentucky: 656–663, 736
Kenya: 1209
Kernohan, Brian J., *see* Seaman, D. Erran
Kesterson Reservoir: 1273–1281
Kestrel, American: 1067, 1273–1281
selenium accumulation and effects on reproduction in captive American kestrels fed selenomethionine, 502–511
Keterson National Wildlife Refuge: 509
Kidney fat index:
adaptive fetal sex allocation in elk: evidence and implications, 1109–1117
Kidneywood, Texas: 570
Killdeer: 270, 1013
Kimball, Bruce A., *see* Mason, J. Russell
Kingbird, eastern: 807, 1013
Kingbird, western: 1013
Kingery, James L., *see* Alpe, Michael J.
Kingfisher, belted: 170–177, 489
Kinglet, golden-crowned: 170–177, 187–188, 484–496
Kinglet, ruby-crowned: 170–177, 188
Kirby, Ronald E., and Glen A. Sargeant. Survival of postfledging mallards in northcentral Minnesota, 403–408
Kirsch, Eileen M., and John G. Sidle. Status of the interior population of least tern, 470–483
Kisatchie National Forest: 825
Kite, Mississippi: 83
Klaas, Erwin E., *see* Hubbard, Michael W.
Klaus, Nathan A., *see* Dettmers, Randy
Klein, David R., *see* Post, Eric
Koa haole: 998
Kochert Michael N., Karen Steenhof, Leslie B. Carpenter, and John M. Marzluff. Effects of fire on golden eagle territory occupancy and reproductive success, 773–780
Kochia: 1300
Kochia scoparia, see Kochia
Kohlmann, Stephen G. Adaptive fetal sex allocation in elk: evidence and implications, 1109–1117
Koloa, *see* Duck, Hawaian
Koneff, Mark D., *see* Williams, Byron K.
Koontz, Fred W., *see* Ostro, Linde E.
Krameria grayi, see Rhatany, white
Krascheninnikovia, lanata, see Winterfat
Krementz, David G., and Jeffery S. Chrisite. Scrub-successional bird community dynamics in young and mature longleaf pine-wiregrass savannahs, 803–814
Krzysik, Anthony J., *see* Duda, Jeffery J.
Kunkel, Kryan, and Daniel H. Pletscher. Species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
———, Toni K. Ruth, Daniel H. Pletscher, and Maurice G. Hornocker. Winter prey selection by wolves and cougars in and near Glacier National Park, Montana, 901–910
Kurta, Charlotte, *see* Sullivan Thomas P.
Kurzejeski, Eric W., *see* McCoy, Timothy D.

L

Labisky, Ronald F., Karl E. Miller, and Christine S. Hartless. Effect of Hurricane Andrew on survival and movements of white-tailed deer in the everglades, 872–879
Labrador: 580–587
Lagopus lagopus scoticus, see Grouse, red
Lagopus lagopus, see Grouse, willow
Lagopus lagopus, see Ptarmigan, willow
Lagopus mutus, see Ptarmagin, Scottish
Lake Woodruff National Wildlife Refuge: 1193–1200
LANDSAT: 1212–1221
Lanius ludovicianus, see Shrike, loggerhead
Lank. David B., *see* Lougheed, Lynn W.

Larch, eastern: 168, 347
Larch, western: 902, 1083
Larix laricina, see Larch, eastern
Larix occidentalis, see Larch, western
Lark, horned: 270–272
Larrea tridentata, see Creosote
Larus glaucescens, see Gull, glaucous-winged
Lasionycteris noctivagans, see Bat, silver-haired
Lasiurus borealis, see Bat, red
Lasiurus cinereus, see Bat, hoary
Lasley, Bill L., *see* Stoops, Monica A.
Laterallus jamaicensis, see Rail, black
Lautenschlager, R. A., William J. Dalton, Marilyn L. Cherry, and Jeri L. Graham. Conifer release alternatives increase aspen forage quality in northwestern Ontario, 1320–1326
Lawrance, Peter S., *see* Flemming, Stephen P.
Lead: 790
Leafloor, James O., *see* Abraham, Kenneth F.
Leberg, Paul L., and Darrell L. Ellsworth. Further evaluation of the genetic consequences of translocation on southeastern white-tailed deer populations, 327–334
Leberg, Paul L., *see* Mallach, Troy J.
Lee, Bruce A. Vander, R. Scott Lutz, Leslie A. Hansen, Nancy E. Matthews. Effects of supplemental prey, vegetation, and time on success of artificial nests, 1299–1305
Legare, Michael L., William R. Eddleman, P. A. Buckley, and Colleen Kelly. The effectiveness of tape playback in estimating black rail density, 116–125
Leif, Anthony P., *see* Gabbert, Andy E.
Leipoa ocellata, see Fowl, mallee
Leks:
land-use patterns surrounding greater prairie-chicken leks in northwestern Minnesota, 189–198
Leopold, Bruce D., *see* Chamberlain, Michael J.
———, *see* Conner, Mike
———, *see* Miller, Darren A.
Lepomis spp., *see* Sunfish
Lepus californicus, see Rabbit, black-tailed jack
Lercel, Barbara A., Richard M. Kaminski, and Robert R. Cox, Jr. Mate loss in winter affects reproduction of mallards, 621–629
Lethal dose: 1138–1151
Leucaena leucocephala, see Koa haole
Leucophyllum frutescens, see Cenizo
Libocedrus decurrens, see Cedar, incense
Lichen: 337, 339–340
Light, sodium-iodide: 1009–1017
Lily, swamp: 877
Lily, white water: 874
Limbo, gumbo: 874
Lincoln National Forest: 60–76
Link, Wiliam A. Modeling patternin collections of parameters, 1017–1027
Linnell, Michael A., Michael R. Conover, and Tim J. Ohashi. Biases in bird strike statistics based on pilot reports, 997–1003
Lion, mountain: 305, 608, 1104
Lipia ligustrina, see Whitebrush
Lipids:
use of total body electrical conductivity to predict northern bobwhite lipid mass, 695–704
Liquidambar styraciflua, see Gum, sweet
***Listeria* spp.:** 207
Lithocarpus densiflorus, see Oak, tan
Livestock: 346–354, 422–425, 593–612, 896, 925, 927, 939–949, 1067, 1078, 1106
Lizard, blunt-nosed leopard: 1188
Lizard, lava: 1060
Lizard, spiny: 1188
Local resource competition hypothesis: 1110
Lockman, David C., *see* Drewien, Roderick C.
Lonchura malacca, see Mannikin
Lonchura punctulata, see Mannikin
Longspur, chestnut-collared: 530–538
Longspur, McCown's: 270–272
Loon, common: 1363–1364
Lophodytes cucullatus, see Merganser, hooded
Lougheed, Lynn W., André Breault, and David B. Lank. Estimating statistical power to evaluate ongoing waterfowl population monitoring, 1359–1369
Louisiana: 137–146, 220, 329, 470–483, 619, 635, 995
Lovell, Charles, *see* Conner, Mike
Loxodonta africana, see Elephant, African
Luttich, Stuart N., *see* Schaefer, James A.
Lutz, R. Scott, *see* Lee, Bruce A. Vander
Lycaon pictus, see Dog, wild
Lynx: 355, 1104
Lynx lynx, see Lynx
Lynx rufus, see Bobcat
Lysiloma latisiliqua, see Tamarind, wild

M

M-44: 939–949
Macaca fascicularis, see Macaque
Macaque: 563
Mace, Richard D., *see* McLellan, Bruce N.
Maclura pomifera, see Osage orange
Madrone: 45, 56, 205
Magnolia virginiana, see Bay, sweet
Mahogany, mountain: 86, 452
Maine: 173, 176, 185, 441–450, 907
Mallach, Troy J., and Paul L. Leberg. Use of dredged material substrates by nesting terns and black skimmers, 137–146
Malus spp., *see* Apple, crab
Manatee, Florida: 253
***Mandrillus leucophaeus*:** 6–7
Mangrove:
evaluation of translocation for the threatened Florida scrub-jay, 833–842
Mangrove, red: 1203

Mangrove, white: 1203
Manitoba: 429, 621
Mankin, Philip C., and Richard E. Warner. A regional model of the eastern cottontail and land use changes in Illinois, 956–963
Mankin, Philip C., *see* Warner, Richard E.
Mannan, R. William, *see* Boal, Clint W.
———, *see* Hall, Linnea S.
———, *see* Meretsky, Vicky J.
Mannikin: 999–1000
Manti-LaSal National Forest: 607
Maple, big-leaf: 45, 56, 485
Maple, mountain: 1307
Maple, red: 168, 181, 1173
Maple, silver: 630–636
Maple, sugar: 462–466, 965, 1307
Maple, vine: 293, 486, 555
Mare's tail: 1300
Margarops fuscatus, see Thrasher, pearly-eyed
Mark Twain National Forest: 185
Marmot: 313
Marmot, yellow-bellied: 777
Marmota flaviventris, see Marmot, yellow-belied
Marmota spp., *see* Marmot
Marshal, Jason P., and Stan Boutin. Power analysis of wolf-moose functional responses, 396–402
Marten, American: 292
Martes americana, see Marten, American
Martes pennanti, see Fischer
Martin, Karl J., and Robert G. Anthony. Movements of northern flying squirrels in different-aged forest stands of western Oregon, 291–297
Maryland: 329, 857
Marzluff, John M., *see* Kochert, Michael N.
Mason, J. Russell, John Hollick, Bruce A. Kimball, and John J. Johnston. Repellency of Deer Away Big Game Repellent® to eastern cottontail rabbits, 309–314
Massachusetts: 965, 1173–1180
Mast: 927
Masters, Ronald E., *see* James, John B.
Matchwood: 233
Matthews, Lindsay R., *see* Waas, Joseph R.
Matthews, Nancy E., *see* Lee, Bruce A. Vander
Mayfield method: 369–373, 382–389, 525
Mazama americana, see Deer, red brocket
Mazama gouazoubira, see Deer, grey brocket
McCollough, Dale R., *see* Sacks, Benjamin N.
McCorquodale, Scott M. Movements, survival, and mortality of black-tailed deer in the Klickitat basin of Washington, 861–871
McCoy, Timothy D., Mark R. Ryan, Eric W. Kurzejeski, and Loren W. Burger, Jr. Conservation Reserve Program: source or sink habitat for grassland birds in Missouri?, 530–538
McGrane, David G., *see* Avery, Michael L.
McKelvey, Kevin S., *see* Rosenberg, Daniel K.
McKinnon, Donald T., and David C. Duncan. Effectiveness of dense nesting cover for increasing duck production in Saskatchewan, 382–389
McLellan, Bruce N., Fred W. Hovey, Richard D. Mace, John G. Woods, Daniel W. Carney, Michael L. Gibeau, Wayne L. Wakkinen, and Wayne F. Kasworm. Rates and causes of grizzly bear mortality in the interior mountains of British Columbia, Alberta, Montana, Washington, and Idaho, 911–920
McMullen, Craig M., *see* Heffelfinger, James R.
Meadowlark, eastern: 530–538, 1012
Meadowlark, western: 530–538, 999–1001
Mednis, Aivars, *see* Blums, Peter
Melanerpes carolinus, see Woodpecker, red-bellied
Melanerpes erythrocephalus, see Woodpecker, red-headed
Meleagris gallopavo intermedia, see Turkey, Rio Grande
Meleagris gallopavo merriami, see Turkey, Merriam's
Meleagris gallopavo sylvestris, see Turkey, eastern wild
Melospiza lincolnii, see Sparrow, Lincoln's
Melospiza melodia, see Sparrow, song
Mephitis mephitis, see Skunk, striped
Meretsky, Vicky J., and R. William Mannan. Supplemental feeding regimes for Egyptian vultures in the Negev Desert, Israel, 107–115
Merganser, hooded: 656
Merlin: 80–81
Merrill, Michael D., Kim A. Chapman, Karen A. Poiani, and Brian Winter. Land-use patterns surrounding greater prairie-chicken leks in northwestern Minnesota, 189–198
Meslow, E. Charles, *see* Swindle, Keith A.
Mesquite: 78, 451–461, 1010
Mesquite, honey: 570–571, 1067, 1292
Metapopulations: 470–483
  use of population viability analysis to evaluate management options for the endangered Lower Keys marsh rabbit, 251–260
Mexico: 127, 451, 1201–1212, 1231, 1240
Michener, Robert, *see* Jacoby, Michael E.
Michigan: 73, 535
Micropterus dolomieui, see Bass, smallmouth
Microtus californicus, see Vole, California
Microtus oeconomus, see Vole, root
Migrants, Neotropical:
  abundance and productivity of forest songbirds in a managed, unfragmented landscape in Vermont, 180–188
Migration:
  molt migrant Canada geese in northern Ontario and western James Bay, 649–655
  performance and accuracy evaluation of small satellite transmitters, 1349–1358
Miller, Darren A., George A. Hurst, and Bruce D. Leopold. Habitat use of eastern wild turkeys in central Mississippi, 210–222
Miller, Karl E., *see* Labisky, Ronald F.

Miller, Michael R., and Wesley E. Newton. Population energetics of northern pintails wintering in the Sacramento Valley, California, 1222–1238
Millspaugh, Joshua J., *see* Seaman, D. Erran
Mimosa biuncifera, see Mimosa, cat's claw
Mimosa, cat's claw: 1067
Mimus polyglottos, see Mockingbird, northern
Mingo Swamp National Wildlife Refuge: 636, 657–663
Minimum convex polygon: 739–747, 845, 853–860, 875, 1183
Mink: 202, 405
Minnesota: 189–198, 389–395, 403–408, 577–578, 635, 889, 972, 1090, 1102–1105
Mirarchi, Ralph E., *see* Sullivan, J. Daniel, Jr.
Mississippi: 210–222, 261–269, 424, 470–483, 613–620, 635, 656–663, 736, 880–888, 972, 1016, 1261–1272, 1278
Mississippi River: 470–483
Missouri: 185, 208, 228, 470–483, 520, 656–663, 695–704, 972, 1287
Missouri River: 470–483
Mitchell, Joseph C., *see* Carter, Shawn L.
Mitchell, Lee R., Lawrence D. Carlile, and C. Ray Chandler. Effects of southern flying squirrels on nest success of red-cocked woodpeckers, 538–545
Mitochondrial DNA: 327–334
Mniotilta varia, see Warbler, black-and-white
Mockingbird, northern: 807, 1000, 1012
Model, Cox proprotional hazards: 404
Model, Fisher's: 1109
Modeling:
an aerial sightability model for estimating ferruginous hawk population size, 85–97
influence of point count length and repeated visits on habitat model performance, 815–823
modeling patternin collections of parameters, 1017–1027
Modeling, bottom-up: 1066
Modeling, top-down: 1066
Mojave Desert: 1181–1192
Mold, water: 1167
Molothrus ater, see Cowbird, brown-headed
Molt:
molt migrant Canada geese in northern Ontario and western James Bay, 649–655
molt of after-hatching-year mourning doves, 223–231
Monkey, black howler:
a geographic information system method for estimation home range size, 748–755
Monkey, patas: 753
Montana: 173, 315–326, 369, 428–429, 470–483, 782, 784, 787–790, 901–929, 950–955, 1082–1093, 1094–1108
Monte Carlo simulation: 1372
estimation of habitat selection for central-place foraging animals, 1028–1038
Monte Vista National Wildlife Refuge: 129, 135
Montevecchi, William A., *see* Whitaker, Darroch M.
Montopoli, George J., *see* Harmata, Alan R.
Moorhen, common: 122
Moose: 94, 347, 567, 580–587, 901–910, 917, 922–923, 954, 1115, 1320–1326
effects of ear-tagging with radiotransmitters on survival of moose calves, 354–358
power analysis of wolf-moose functional responses, 396–402
species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
Moosehorn National Wildlife Refuge: 185
Morgan, David R., *see* Henderson, Ray J.
Mörschel, Frank M. Use of climatic data to model the presence of oestrid flies in caribou herds, 588–593
Mortality: 1094–1108, 1273–1281
effects of saline drinking water on early gosling development, 990–996
mortality causes and survival rates of hunted and unhunted willow grouse, 722–730
northern bobwhite chick mortality caused by red imported fire ants, 1291–1298
rates and causes of grizzly bear mortality in the interior mountains of British Columbia, Alberta, Montana, Washington, and Idaho, 911–920
survival and cause-specific mortality of adult raccoons in central Mississippi, 880–888
survival and cause-specific mortality of adult bobcats in central Mississippi, 613–620
the efficacy of baits containing 1080 for control of brushtail possums, 1138–1151
Mosillo, Maia, Edward J. Heske, and John D. Thompson. Survival and movements of translocated raccoons in northcentral Illinois, 278–286
Mosley, Jeffrey C., *see* Alpe, Michael J.
Moss, peat: 854
Mouse, cactus: 1075
Mouse, deer: 15, 270, 275, 931, 1067, 1071–1072, 1075, 1077, 1278
relations of small mammal populations to even-aged shelterwood systems: a comment, 1376–1380
relations of small mammal populations to even-aged shelterwood systems: a reply, 1381–1389
Mouse, hispid pocket: 1067, 1071–1072, 1075, 1077
Mouse, house: 931, 1278
Mouse, northern grasshopper: 270, 275, 1067, 1070–1072, 1075, 1077
Mouse, plains harvest: 1067, 1071–1072, 1075, 1077
Mouse, plains pocket: 1067, 1071–1072, 1075, 1077
Mouse, San Joaquin pocket: 931
Mouse, silky pocket: 1067, 1070–1072, 1075, 1077
Mouse, western harvest: 1067, 1071–1072, 1075, 1077, 1278
Mouse, white-footed: 1067, 1071–1072, 1075, 1077, 1175
Mowing: 519, 693, 859

Muddy Creek Wildlife Management Area: 1281–1290
Mueller, James M., C. Brad Dabbert, Stephen Demarais, and Andrew R. Forbes. Northern bobwhite chick mortality caused by red imported fire ants, 1291–1298
Muhlenbergia filipes, see Grass, muhly
Muhlenbergia spp., *see* Muhly
Muhly: 1067
Mullein: 347
Mumme, Ronald L., and Theodore H. Below. Evaluation of translocation for the threatened Florida scrub-jay, 833–842
Murrelet, ancient:
norway rats as predators of burrow-nesting seabirds: insights from stable isotope analyses, 14–25
Mus musculus, see Mouse, house
Muscids: 362
Muskrat: 313
Mussel, California: 1066
Mustard, tumble: 271
Mustela frenata, see Weasel, long-tailed
Mustela nigripes, see Ferret, black-footed
Mustela spp., *see* Weasel
Mustela vison, see Mink
Myiarchus crinitus, see Flycatcher, great-crested
Myna, common: 1000
Myocastor coypus, see Coypu
Myocastor coypus, see Nutria
Myotis, California: 555–561
Myotis californicus, see Myotis, California
Myotis evotis, see Myotis, long-eared
Myotis, fringed: 553–561
Myotis leibii, see Bat, eastern small-footed
Myotis, long-eared: 553–561
Myotis, long-legged: 553–561, 1315
Myotis lucifugus, see Bat, little brown
Myotis septentrionalis, see Bat, northern long-eared
Myotis thysanodes, see Myotis, fringed
Myotis volans, see Myotis, long-legged
Myotis, Yuma: 555–561
Myotis Yumanensis, see Myotis, Yuma
Myrica cerifera, see Myrtle, wax
Myrtle, wax: 117, 874
Mytilus californicus, see Mussel, California
Mytilus spp: 17

N

National Elk Refuge: 130, 1120–1132, 1327
National Elk Refuge Herd: 1123–1132
National Environmental Policy Act: 950
National Oceanic and Atmospheric Administration: 999
Neale, Jennifer C. C., *see* Sacks, Benjamin N.
Nebraska: 470–483, 950–955
Needlerush: 117
Nematodes: 362
Neotoma fuscipes, see Rat, wood
Neotoma micropus, see Rat, southern plains wood
NEPA, *see* National Environmental Policy Act
Nest, artificial:
effects of supplemental prey, vegetation, and time on success of artificial nests, 1299–1305
predation of artificial ground nests on white-tailed prairie dog colonies, 270–277
Nest, cavity:
natural-cavity use by nesting wood ducks in Illinois, 630–638
Net, mist: 805
New England: 369, 1172–1180
New Hampshire: 185
New Mexico: 36–43, 60–76, 104, 126–136, 208, 451, 470–483, 921–929, 933
New York: 635, 972, 1001, 1287
New Zealand: 306, 1056, 1138–1151, 1152–1162
Newfoundland: 167–179
Newt, eastern: 444, 448
Newt, red-spotted: 1175
Newton, Wesley E., *see* Miller, Michael
Nicaragua: 451
Ninebark: 346–354
NOAA, *see* National Oceanic and Atmospheric Administration
Norman, Gary W., *see* Pack, James C.
North American Waterfowl Management Plan: 382–383
evaluation of waterfowl conservation under the North American Waterfowl Management Plan, 417–440
North Carolina: 143, 228, 329, 843–852
North Dakota: 428–429, 470–483, 531, 534–536, 995, 1077
Norway: 728
Notonectids: 362
Notophthalamus viridescens, see Newt, red-spotted
Notophthalmus viridescens, see Newt, eastern
Noxubee National Wildlife Refuge: 657–663, 1261–1272
Nuthatch, brown-headed: 807
Nuthatch, red-breasted: 170–177, 188, 489
Nuthatch, white-breasted: 188
Nutria: 313
local determinants of coypu distribution along the Luján River, eastcentral Argentina, 895–900
Nycticorax nycticorax, see Heron, black-crowned night
Nymphaea odorata, see Lily, white water
Nyssa sylvatica, see Gum, black

O

Oak: 32, 78, 212, 451–461, 927
residual effects of thinning and high white-tailed deer densities on northern redback salamanders in southern New england oak forests, 1172–1180
Oak, blackjack: 1010
Oak, bur: 199, 731

Oak, California black: 45, 56
Oak, California live: 45
Oak, chestnut: 965
Oak, emory: 456
Oak, Gambel's: 36–37, 316, 927
Oak, live: 874, 1292
Oak, net-leaf:451–461
Oak, northern red: 181, 199, 731, 965, 1173
Oak, Oregon white: 45, 205, 862–863
Oak, post: 1010, 1282
Oak, sand shinnery: 1010, 1067
Oak, silver-leaf: 451–461
Oak, southern red: 1282
Oak, tan: 45, 56–57
Oak, water: 874
Oak, white: 199, 731, 965, 1282
Oakleaf, Bob, *see* Harmata, Alan R.
Oats: 956–963
Oceanodroma furcata, see Petrel, fork-tailed storm
Oceanodroma leucorhoa, see Petrel, Leach's storm
Oceanspray: 347
Ocotillo: 452, 1183
Odocoileus hemionus columbianus, see Deer, Columbian black-tailed
Odocoileus hemionus, see Deer, Mule
Odocoileus virginianus, see Deer, white-tailed
Odocoileus virginianus borealis, see Deer, white-tailed
Odocoileus virginianus seminolus, see Deer, white-tailed
Ohashi, Tim J., *see* Linnell, Michael A.
Okapi: 567
Okapia johnstoni, see Okapi
Oklahoma: 270, 470–483, 1009–1017
Olding, Ronald J., *see* Heffelfinger, James R.
Oligochaetes: 362
Olsen, Steven C., *see* Roffe, Thomas J.
Onchorhynchus clarki, see Trout, cutthroat
Onchorhynchus spp., *see* Salmon
Oncorhynchus mykiss, see Trout, rainbow
Oncorhynchus nerka, see Salmon, sockeye
Oncorhynchus tshawytscha, see Salmon, chinook
Ondatra zibethica, see Muskrat
Ontario: 283, 649–655, 1306–1326
Onychomys leucogaster, see Mouse, northern grasshopper
Oporornis formosus, see Warbler, Kentucky
Oporornis philadelphia, see Warbler, mourning
Oporornis tolmiei, see Warbler, MacGillivray's
Opossum: 278, 370, 687, 1300
Opuntia lindheimeri, see Cactus, pricklypear
Opuntia ramosissima, see Cactus, pencil cholla
Opuntia spp., *see* Cacti
Oregon: 57, 204–210, 291–297, 346–347, 467, 484–496, 553–561, 784, 794–802, 1109–1117, 1212–1221, 1239–1240
Oregon Coast Range: 180, 484–496, 553–561
Organophosphates: 81, 786, 790
Oriole, orchard: 807–808
Oryctolagus cuniculus, see Rabbit
Oryzomys argentatus, see Rat, silver rice
Oryzopis contracta, see Grass, Indian rice
Osage orange: 687
Osborne, David A., *see* Frawley, Brian J.
Osprey: 93, 795
Ostracods: 362
Ostro, Linde E. T., Truman P. Young, Scott C. Silver, Fred W. Koontz. A geographic information system method for estimating home range size, 748–755
Otis, David L., and Gary C. White. Autocorrelation of location estimates and the analysis of radiotracking data, 1039–1044
Otter, sea: 794, 1066
Otus asio, see Owl, eastern screech
Ouray National Wildlife Refuge: 133, 135
Ovenbird: 170–177, 183–188, 807
Ovis canadensis canadensis, see Sheep, Rocky Mountain bighorn
Owl, barn: 57, 64, 507, 999–1000, 1067, 1273–1281
Owl, barred: 202
Owl, burrowing: 1067, 1273–1281
Owl, California spotted: 36, 41, 45, 56, 1188
Owl, eastern screech: 80–81, 508–509, 807, 1274–1281
Owl, great-horned: 57, 64, 81, 202, 292, 405, 733, 1274–1281
Owl, Mexican spotted:
effects of helicopter noise on Mexican spotted owls, 60–76
habitat composition and configuration around Mexican spotted owl nest and roost sites in the Tularosa Mountains, New Mexico, 36–43
Owl, northern hawk: 57
Owl, northern spotted: 36, 39, 41–42, 291, 457, 1188
estimation of habitat selection for central-place foraging animals, 1028–1038
forest stand characteristics and reproduction of northern spotted owls in managed north-coastal California forests, 44–59
old-forest distribution around spotted owl nests in the central Cascade Mountains, Oregon, 1212–1221
Owl, short-eared: 999–1001
response of northern harriers and short-eared owls to grassland management in Illinois, 517–523
Owl, tawny: 57
Oxyura jamaicensis, see Duck, ruddy

P

Pac, David F., *see* Unsworth, James W.
Paca: 2–3
Pace, Richard M., and Alan D. Afton. Direct recovery rates of lesser scaup banded in northwest Minnesota: sources of heterogeneity, 389–395
Pack, James C., Gary W. Norman, Curtis I. Taylor, David E. Steffen, David A. Swanson, Kenneth H. Pollock, Russell Alpizar-Jara. Effects of fall hunt-

ing on wild turkey populations in Virginia and West Virginia, 964–975
Palm, cabbage: 874
Palmer, Mitchell V., *see* Roffe, Thomas J.
Pan traglodytes, see Chimpanzee
Panama: 546
Pandion haliaetus, see Osprey
Panicum dichotomiflorum, see Panicum, fall
Panicum, fall: 1261–1272
Panicum hemitomon, see Cane, maiden
Panicum maximum, see Grass, Guinea
Panicum spp., *see* Grass, panic
Panicum virgatum, see Grass, switch
Panther, Florida: 305, 878
Panthera onca, see Jaguar
Panthera tigris, see Tiger
***Papio cynocephalus*:** 6–7
Papua New Guinea: 1193
Parabuteo unicinctus, see Hawk, Harris'
Parameters:
modeling patternin collections of parameters, 1017–1027
Paroaria coronata, see Cardinal, red-crested
Partridge: 705, 723
Partridge, gray: 705
effects of radiotransmitters on survival and reproductive success of gray partridge, 1044–1051
Parula americana, see Parula, northern
Parula, northern: 807
Parus atricapillus, see Chickadee, black-capped
Parus, hudsonicus, see Chickadee, boreal
Paspalum: 141
Paspalum spp., *see* Paspalum
Passer domesticus, see Sparrow, house
Passerculus sandwichensis, see Sparrow, savannah
Passerella iliaca, see Sparrow, fox
Passerina ciris, see Bunting, painted
Pater, Larry L., *see* Delany, David K.
Pear, flowering: 309–314
Peccary, collared: 3–5
seasonal variation of home range of collared peccary in tropical rain forests of French Guiana, 546–552
Peccary, white-lipped: 3–5
Peeper, northern spring: 1175
Peery, M. Zachariah, R. J. Gutiérrez, and Mark E. Seamans. habitat composition and configuration around Mexican spotted owl nest and roost sites in the Tularosa Mountains, New Mexico, 36–43
Peewee, eastern: 183–188, 807–809
Peewee, western: 489–490
Penguin, Galápagos: 1058
Pennsylvania: 104, 329, 857
Penteriani, Vincenzo. Dawn and morning goshawk courtship vocalizations as a method for detecting nest sites, 511–516
Perca flavescens, see Perch, yellow
Perch, yellow: 1166
Percival, H. Franklin, *see* Rice, Kenneth G.
Perdix perdix, see Partridge, gray
Perisoreus canadensis, see Jay, gray
Perognathus favescens, see Mouse, plains pocket
Perognathus flavus, see Mouse, silky pocket
Perognathus inornatus, see Mouse, San Joaquin pocket
Peromyscus eremiscus, see Mouse, cactus
Peromyscus leucopus, see Mouse, white-footed
Peromyscus maniculatus, see Mouse, deer
Persea palustris, see Bay, red
Persimmon, Mexican: 571
Peru: 3–13, 1209
Petrel, dark-rumped: 1058
Petrel, fork-tailed storm: 15
Petrel, Leach's storm: 15
***Petrodomus tetradactylus*:** 6–7
Phasianus colchicus, see Pheasant, ring-necked
Pheasant, ring-necked: 200, 777, 957, 999–1001
declining survival of ring-necked pheasant chicks in Illinois during the late 1900s, 705–710
site selection and nest success of ring-necked pheasants as a function of location in Iowa landscapes, 976–989
survival of ring-necked pheasant hens during spring in relation to landscape features, 147–154
survival and habitat use by ring-necked pheasants during two diparate winters in South Dakota, 711–722
Pheucticus ludovicianus, see Grosbeak, rose-breasted
Pheuticus melanocephalus, see Grosbeak, black-headed
Philippines: 6
Philo, L. Michael, *see* Roffe, Thomas J.
Philopatry: 332
Phleum pratense, see Timothy
Phlox: 272
Phlox hoodii, see Phlox
Phoenicopterus chilensis, see Flamingo, Chilean
Phoenicopterus ruber ruber, see Flamingo, American
Phragmites australis, see Reed, giant
Physocarpus malvaceus, see Ninebark
Picea abies, see Spruce, Norway
Picea engelmannii, see Spruce, Engelmann
Picea glauca, see Spruce, white
Picea mariana, see Spruce, black
Picea rubens, see Spruce, red
Picea stichensis, see Spruce, Sitka
Pickerelweed: 874
Picoides arcticus, see Woodpecker, black-bellied
Picoides borealis, see Woodpecker, red-cockaded
Picoides pubescens, see Woodpecker, downy
Picoides villosus, see Woodpecker, hairy
Pierce, D. John, *see* Watson, James W.
Pig, feral:
range expansion, population sizes, and management of wild pigs in California, 298–308
Pigeon, band-tailed: 229, 490
Pigeon, feral: 81, 229
Pigeon, white-crowned: 242, 258

***Piliocolobus pennanti*:** 6–7
Pils, Andrew C., Robert A. Garrot, and John J. Borkowski. Sampling and statistical analysis of snow-urine allantoin:cretinine ratios, 1118–1132
Pine, Chihuahua: 452, 456
Pine, eastern white: 181
Pine, engelmann: 452
Pine, jack: 462, 468, 1307
Pine, Jeffery: 45
Pine, loblolly: 26, 211, 261, 613, 804, 825, 844, 846, 881
Pine, lodgepole: 317, 902, 1083, 1119–1120, 1377, 1382
Pine, longleaf: 26–27, 32–33, 261, 539
scrub-successional bird community dynamics in young and mature longleaf pine-wiregrass savannahs, 803–814
selection of pines for foraging by red-cockaded woodpeckers, 843–852
Pine, pinyon: 37, 316
Pine, pond: 844, 846
Pine, ponderosa: 36–37, 45, 61, 99, 347, 795, 863
Pine, red: 462, 1307
Pine, sand: 26
Pine, scotch: 355
Pine, shortleaf: 261, 825, 846, 1282
Pine, slash: 26–35, 804, 844, 846
Pine, southwestern white: 36, 61
Pine, sugar: 45
Pine, western white: 347
Pine, white: 456, 462, 816, 1173, 1306–1319, 1321
Pinicola enucleator, see Grosbeak, pine
Pintail, northern: 383–384, 417–440, 647, 1258, 1359–1369
population energetics of northern pintails wintering in the Sacramento Valley, California, 1222–1238
Pinus avacahuite, see Pine, white
Pinus banksiana, see Pine, jack
***Pinus cemboides*:** 452
Pinus clausa, see Pine, sand
Pinus contorta, see Pine, lodgepole
Pinus echinata, see Pine, shortleaf
Pinus edulis, see Pine, pinyon
Pinus elliottii, see Pine, slash
Pinus engelmanii, see Pine, engelmann
Pinus jeffeyi, see Pine, Jeffery
Pinus lambertiana, see Pine, sugar
Pinus leiophylla, see Pine, Chihuahua
Pinus monticola, see Pine, western white
Pinus palustris, see Pine, longleaf
Pinus ponderosa, see Pine, ponderosa
Pinus resinosa, see Pine, red
Pinus serotina, see Pine, pond
Pinus strobiformis, see Pine, southwestern white
Pinus strobus, see Pine, eastern white
Pinus strobus, see Pine, white
Pinus sylvestris, see Pine, scotch
Pinus taeda, see Pine, loblolly
Pipilo erythrophthalmus, see Towhee, eastern
Pipilo maculatus, see Towhee, spotted
Piranga ludoviciana, see Tanager, western
Piranga olivacea, see Tanager, scarlet
Piranga rubra, see Tanager, summer
Pisaster ochraceus, see Sea star
Pisenti, Jacqueline M., *see* Santolo, Gary M.
Pityopsis graminifolia: 29
Platanus occidentalis, see Sycamore, American
Platanus sargentii, see Cottonwood, plains
Platanus wrightii, see Sycamore
Platte River: 470–483
Plethodon cinereus, see Salamander, northern redback
Pletscher, Daniel H., *see* Boyd, Diane K.
———, *see* Kunkel, Kryan E.
Plover, lesser Pacific golden: 999–1001
Plover, mountain: 270
Plowman, Bruce W., *see* Conner, Mike
———, *see* Chamberlain, Michael J.
Pluvialis fulva, see Plover, lesser Pacific golden
Pneumonia: 719
Poa secunda, see Grass, Sandberg blue
Poa spp., *see* Grass, blue
Pochard, common: 657
Poecile carolinensis, see Chickadee, Carolina
Poecile rufescens, see Chickadee, chestnut-backed
Poiani, Karen A., *see* Merrill, Michael D.
Point Reyes National Seashore: 561–569
Polioptila caerulea, see Gnatcatcher, blue-gray
Pollock, Kenneth H., *see* Pack, James C.
———, *see* Tsai, Kuenhi
Polygonum hydropiperoides, see Smartweed, swamp
Polygonum spp., *see* Smartweed
Polystichum munitum, see Fern, sword
Pontederia lanceolata, see Pickerelweed
Pooecetes gramineus, see Sparrow, vesper
Poplar, balsam: 575
Population dynamics:
population dynamics of Zenaida doves in Cidra, Puerto Rico, 232–244
Population estimate:
an aerial sightability model for estimating ferruginous hawk population size, 85–97
Population viability analysis:
use of population viability analysis to evaluate management options for the endangered Lower Keys marsh rabbit, 251–260
Populus spp., *see* Cottonwood
Populus balsamifera, see Popular, balsam
Populus deltoides, see Cottonwood, eastern
Populus fremontii, see Cottonwood, Fremont
Populus tremuloides, see Aspen, quaking
Portulaca oleracae, see Purslane
Porzana carolina, see Sora
Porzana tabuensis, see Crakes, spotless
Possum, brushtail:
the efficacy of baits containing 1080 for control of brushtail possums, 1138–1151
Possum, mountain brushtail: 253
Post, Eric, and David R. Klein. Caribou calf produc-

tion and seasonal range quality during a population decline, 335–345
Potentilla palustris, see Cinquefoil, marsh
Power analysis: 1060
estimating statistical power to evaluate ongoing waterfowl population monitoring, 1359–1369
power analysis of wolf-moose functional responses, 396–402
the insignificance of statistical significance testing, 763–772
Power lines: 128, 441, 728
Prairie: 417–440, 523–538, 639–649, 1066–1081
Prairie dog, black-tailed: 275–276
Prairie dog, white-tailed:
predation of artificial ground nests on white-tailed prairie dog colonies, 270–277
Prairie Pothole Region: 276, 382, 421–440, 1078
Precipitation:
influence of precipitation timing and summer temperatures on reproduction of Gambel's quail, 154–161
Predation, density-dependent: 396–402
Predation:
effect of preventive coyote hunting on sheep losses to coyote predation, 606–612
effects of southern flying squirrels on nest success of red-cockaded woodpeckers, 538–545
effects of supplemental prey, vegetation, and time on success of artificial nests, 1299–1305
factors influencing wild turkey hen survival in southcentral Iowa, 731–738
mortality causes and survival rates of hunted and unhunted willow grouse, 722–730
natural-cavity use by nesting wood ducks in Illinois, 630–638
norway rats as predators of burrow-nesting seabirds: insights from stable isotope analyses, 14–25
predation and survival of white-tailed deer fawns in northcentral New Brunswick, 574–579
predation of artificial ground nests on white-tailed prairie dog colonies, 270–277
relative vulnerability of coyotes to removal methods on a northern California ranch, 939–949
reproduction in a declining population of wild turkeys in Arkansas, 1281–1290
species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
territoriality and breeding status of coyotes relative to sheep predation, 593–605
Prescribed burning: 519, 527, 535, 538, 693, 803, 840, 859
Prey selection:
winter prey selection by wolves and cougars in and near Glacier National Park, Montana, 901–910
Prey, supplemental:
effects of supplemental prey, vegetation, and time on success of artificial nests, 1299–1305
Procyon lotor, see Raccoon
Progestrone: 564–561
Prosopis glandulosa, see Mesquite, honey
Prosopis spp., *see* Mesquite
Prosopium williamsonii, see Whitefish, mountain
Protein, crude: 346–354
Prunus pensylvanica, see Cherry, pin
Prunus serotina, see Cherry, black
Pseudotsuga menziesii, see Fir, Douglas
Psuedotsuga macrocarpa, see Fir, big-cone
Ptarmigan, Scottish: 1287
Ptarmigan, willow: 163, 165
Pteridium aquillinum, see Fern, bracken
Pterodroma phaeopygia, see Petrel, dark-rumped
Ptychoramphus aleuticus, see Auklet, Cassin's
Pueo, *see* Owl, short-eared
Puerto Rico: 232–244
Puffinus newelli, see Shearwater, Newell's
Puma concolor, see Cougar
Puma concolor, see Lion, mountain
Purshia tridentata, see Bitterbush, antelope
Purslane: 141
Purvis, Jon R., *see* Gabbert, Andy E.
PVA, *see* Population viability analysis
Pyracantha coccinea, see Firebush
Pyrrhula pyrrhula, see Finch, bull
Pyrus calleryana, see Pear, flowering

Q

Quail, Gambel's: 670
influence of precipitation timing and summer temperatures on reproduction of Gambel's quail, 154–161
Quail, Japanese: 273, 508
Quail, northern bobwhite: 159, 723, 807, 1007, 1287
energy-based carrying capacity for quails, 664–674
macrohabitat effects on summer survival, movements, and clutch success of northern bobwhite in Kansas, 675–685
microhabitat selection by nesting and brood-rearing northern bobwhite in Kansas, 686–694
northern bobwhite chick mortality caused by red imported fire ants, 1291–1298
passive integrated transponders and patagial tag markers for northern bobwhite chicks, 162–166
slack in the configuration of habitat patches for northern bobwhites, 245–250
use of total body electrical conductivity to predict northern bobwhite lipid mass, 695–704
Quail, scaled: 1287
Quasi-likelihood:
modeling patternin collections of parameters, 1017–1027
Quebec: 176, 603
Quercus alba, see Oak, white
Quercus chrysolepis, see Oak, California live
Quercus emoryi, see Oak, enory
Quercus falcata, see Oak, southern red
Quercus gambelii, see Oak, Gambel's

Quercus garryana, see Oak, Oregon white
Quercus harvardii, see Oak, sand shinnery
Quercus hypoleucoides, see Oak, silver-leaf
Quercus kelloggii, see Oak, California black
Quercus macrocarpa, see Oak, bur
Quercus marilandica, see Oak, blackjack
Quercus nigra, see Oak, water
Quercus prinus, see Oak, chestnut
Quercus rubra, see Oak, northern red
Quercus rugosa, see Oak, net-leaf
Quercus spp., *see* Oak
Quercus stellata, see Oak, post
Quercus virginiana, see Oak, live
Quiscalus major, see Grackle, boat-tailed
Quiscula quiscula, see Grackle, common

R

Rabbit: 298
Rabbit, black-tailed jack: 773, 777–778, 931, 1066–1081, 1278
Rabbit, desert cottontail: 777, 931, 1066–1081, 1278
Rabbit, eastern cottontail:
a regional model of the eastern cottontail and land use changes in Illinois, 956–963
repellency of Deer Away Big Game Repellent® to eastern cottontail rabbits, 309–314
Rabbit, Lower Keys marsh:
use of population viability analysis to evaluate management options for the endangered Lower Keys marsh rabbit, 251–260
Rabbit, New England cottontail: 258
Rabbitbrush: 86
Rabies: 278
Raccoon: 257, 370, 524, 615, 630–636, 687, 736, 877, 985, 1286, 1300
survival and cause-specific mortality of adult raccoons in central Mississippi, 880–888
survival and movements of translocated raccoons in northcentral Illinois, 278–286
survivorship of a nonharvested raccoon population in south Texas, 889–894
Radiotelemetry: 116–125, 147–154, 189–222, 253–258, 262–269, 278–286, 293–297, 396–402, 451–461, 582–585, 595–603, 614–619, 640–647, 686–694, 711–738, 781–793, 853–894, 911–920, 939–949, 964–989, 1084–1091, 1096–1108, 1119–1132, 1181–1192, 1282–1290
autocorrelation of location estimates and the analysis of radiotraking data, 1039–1044
effects of ear-tagging with radiotransmitters on survival of moose calves, 354–358
effects of radiotransmitters on survival and reproductive success of gray partridge, 1044–1051
effects of violation of assumptions for survival analysis methods in radiotelemetry studies, 1369–1375
performance and accuracy evaluation of small satellite transmitters, 1349–1358
seasonal movements of sandhill cranes radiomarked in Yellowstone National Park and Jackson Hole, Wyoming, 126–136
Raedeke, Kenneth J., *see* Seaman, D. Erran
Ragwort, golden: 1328
Rail, black:
the effectiveness of tape playback in estimating black rail density, 116–125
Rail, king: 122
Rail, Virginia: 122
Rail, Yuma clapper: 122
Rainforest:
seasonal variation of home range of collared peccary in tropical rain forests of French Guiana, 546–552
Rallus elegans, see Rail, king
Rallus limicola, see Rail, Virginia
Rallus longirostris yumanensis, see Rail, Yuma clapper
Rana aurora, see Frog, red-legged
Rana catesbeiana, see Frog, bull
Rana palustris, see Frog, pickerel
Rana sylvatica, see Frog, wood
Random effects:
modeling patternin collections of parameters, 1017–1027
Rangifer tarandus caribou, see Caribou, woodland
Rangifer tarandus, see Caribou
Raspberry: 443, 1320–1326
Rat, black: 15, 1058
Rat, hispid cotton: 1067, 1071–1072, 1075, 1077
Rat, kangaroo: 931
Rat, Norway: 931
norway rats as predators of burrow-nesting seabirds: insights from stable isotope analyses, 14–25
Rat, Ord's kangaroo: 1066–1081
Rat, roof: 242
Rat, silver rice: 258
Rat, southern plains wood: 1067, 1070–1072, 1075, 1077
Rat, wood: 44, 48, 57
Rattlesnake, eastern diamondback: 256
Rattus rattus, see Rat, black
Rattus rattus, see Rat, roof
Raven, American: 1300
Raven, Chihuahuan: 1067, 1069
Raven, common: 489, 864, 1300
RB51:
biosafety of parenteral *Brucella abortus* RB51 vaccine in bison calves, 950–955
Recruitment: 580–587
Recurvirostra americana, see Avocet, American
Red River Valley: 189–198
Red Wine Mountain Caribou Herd: 580–587
Redhead: 393, 1363–1364
Redstart, American: 184–188, 807
Reed, giant: 1194, 1196
Regulus calendula, see Kinglet, ruby-crowned
Regulus satrapa, see Kinglet, golden-crowned

Reindeer, *see* Caribou
Reintroduction:
evaluation of translocation for the threatened Florida scrub-jay, 833–842
reintroduction and postrelease movements of red-cockaded woodpecker groups in eastern Texas, 824–832
Reiser, M. Hildegard, *see* Delany, David K.
Reithrodontomys megalotis, see Mouse, western harvest
Reithrodontomys montanus, see Mouse, plains harvest
Reitz, François, *see* Bro, Elisabeth
Repellent:
effectiveness of Hot Sauce® and Deer Away® repellents for deterring elk browsing of aspen sprouts, 1327–1336 plant growth regulator (Stronghold™) enhances repellency of anthraquinone formulation (Flight Control™) to Canada geese, 1336–1343
repellency of Deer Away Big Game Repellent® to eastern cottontail rabbits, 309–314
repellents to reduce cable gnawing by northern pocket gophers, 1344–1349
Reproduction: 1094–1108
adaptive fetal sex allocation in elk: evidence and implications, 1109–1117
forest stand characteristics and reproduction of northern spotted owls in managed north-coastal California forests, 44–59
effects of radiotransmitters on survival and reproductive success of gray partridge, 1044–1051
evaluation of waterfowl conservation under the North American Waterfowl Management Plan, 417–440
influence of precipitation timing and summer temperatures on reproduction of Gambel's quail, 154–161
mate loss in winter affects reproduction of mallards, 621–629
microhabitat selection by nesting and brood-rearing northern bobwhite in Kansas, 686–694
reproduction and survival of Rio Grande turkeys in Oregon, 204–210
reproduction in a declining population of wild turkeys in Arkansas, 1281–1290
selenium accumulation and effects on reproduction in captive American kestrels fed selenomethionine, 502–511
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
use of fecal steroid metabolites to estimate the pregnancy rate of a free-ranging herd of tule elk, 561–569
wood duck brood movements and habitat use on prairie rivers in South Dakota, 639–649
Restani, Marco, *see* Harmata, Alan R.
Restocking: 327–334
Retrospective analysis: 1359–1369
Rhatany, white: 1183
Rhinoceros, black: 567
Rhizophora mangle, see Mangrove
Rhizophora mangle, see Mangrove, red
Rhododendron macrophyllum, see Rhododendron, Pacific
Rhododendron, Pacific: 293
Rhyan, Jack C., *see* Roffe, Thomas J.
***Rhynchocyon chrysopygus*:** 6–7
Rhynchospora globularis, see Rush, beak
Rhyncops niger, see Skimmer, black
Ribes viscosissimum, see Currant, sticky
Rice: 427, 1003–1008, 1247
Rice, Kenneth G., H. Franklin Percival, Allan R. Woodard, and Michael L. Jennings. Effects of egg and hatchling harvest on American alligators in Florida, 1193–1200
Richardson, David M., *see* Blums, Peter
***Rinorea* spp.:** 547
Ripple, William J., *see* Swindle, Keith A.
Rivera-Milan, Frank F. Population dynamics of zenaida doves in Cidra, Puerto Rico, 232–244
Roads: 441
Robbins, Charles T., *see* Jacoby, Michael E.
Robel, Robert J., *see* Hughes, John P.
Robin, American: 170–177, 187–188, 490
Robinson, John G., and Richard E. Bodmer. Towards wildlife management in tropical forests, 1–13—Invited paper
Rocke, Tonie E., and Michael D. Samuel. Water and sediment characterics associated with avian botulism outbreaks in wetland, 1249–1260
———, Ned H. Euliss, Jr., and Michael D. Samuel. Enviromental characteristics associated with the occurrence of avian botulism in wetlands of a northern California refuge, 358–368
Rocky Mountain National Park: 1327
Roffe, Thomas J., Jack C. Rhyan, Keith Aune, L. Michael Philo, Darl R. Ewalt, Thomas Gidlewski, Steven G. Hennager. Brucellosis in Yellowstone National Park bison: quantitative serology and infection, 1132–1137
———, Steven C. Olsen, Thomas Gidlewski, Allen E. Jensen, Mitchell V. Palmer, and Royce Huber. Biosafety of parenteral *Brucella abortus* RB51 vaccine in bison calves, 950–955
Roost sites: 417–440, 1315
habitat composition and configuration around Mexican spotted owl nest and roost sites in the Tularosa Mountains, New Mexico, 36–43
Rosa spp., *see* Rose
Rosa woodsii, see Rose, wild
Rose: 346–354
Rose, multiflora: 310
Rose, wild: 1328
Rose multiflora, see Rose, multiflora
Rose-apple: 233
Rosenberg, Daniel K., and Kevin S. McKelvey. Esti-

mation of habitat selection for central-place foraging animals, 1028–1038
Rubus idaeus, see Raspberry
Rubus parviflorus, see Thimbleberry
Rubus spectabilis, see Salmonberry
Rudolph, D. Craig, *see* Carrie, N. Ross
Rue, meadow: 1328
Ruppia maritima, see Grass, widgeon
Rusch, Donald H., *see* Abraham, Kenneth F.
———, *see* Taylor, J. Scott
Rush, beak: 874, 1261–1272
Rush, spike: 874
Ruth, Toni K., *see* Kunkel, Kryan E.
Ryan, Mark R., *see* McCoy, Timothy D.

S

Sabal palmeatto, see Palm, cabbage
Sabine National Forest: 824–832
Sacks, Benjamin N., Karen M. Blejwas, and Michael M. Jaeger. Relative vulnerability of coyotes to removal methods on a northern California ranch, 939–949
Sacks, Benjamin N., Michael M. Jaeger, Jennifer C. C. Neale, Dale R. McCullough. Territoriality and breeding status of coyotes relative to sheep predation, 593–605
Sacobatus vermiculatus, see Greasewood
Sacramento National Wildlife Refuge: 358–368, 1250–1260
Sage, birdsfoot: 272
Sagebrush: 86–87, 1119
Sagebrush, big: 271, 316–317, 774, 1119
Sagebrush, Wyoming big: 86–87
Sagittaria lancifolia, see Flag
Salal: 16–17, 21–22, 293, 486
Salamander, four-toed: 1175
Salamander, mole: 1172
Salamander, northern dusky: 1175
Salamander, northern redback: 444, 448
residual effects of thinning and high white-tailed deer densities on northern redback salamanders in southern New england oak forests, 1172–1180
Salamander, northern two-lined: 1175
Salamander, spotted: 441–450, 1175
Salinity: 1253–1257
effects of saline drinking water on early gosling development, 990–996
Salix arctica, see Willow, Arctic
Salix gooddingii, see Willow, Gooding's
Salix scouleriana, see Willow, Scouler
Salix spp., *see* Willow
Salmon: 791, 912, 916, 921–929
Salmon, chinook: 924
Salmon, sockeye: 924
Salmonberry: 485
Salsola kali, see Thistle, Russian
Saltbush, Gardner: 86, 141, 271–272
Sambucus spp., *see* Elderberry
Sample size: 1039–1044
effects of sample size on kernel home range estimates, 739–747
Samuel, Michael D., *see* Rocke, Tonie E.
Sandegren, Finn, *see* Swenson, Jon E.
Sandpiper, spotted: 170–177
Sanganois Conservation Area: 630–636
Santolo, Gary M., Julie T. Yamamoto, Jacqueline M. Pisenti, Barry W. Wilson. Selenium accumulation and effects on reproduction in captive american kestrels fed selenomethionine, 502–511
———. Selenium in blood of predatory birds from Kesterson Reservoir and other areas in California, 1273–1281
Saprolegnia ferax, see Mold, water
Sapsucker, red-breasted: 489–490
Sapsucker, yellow-bellied: 188
Sargeant, Glen A., *see* Kirby, Ronald E.
Saskatchewan: 382–389
Savannah:
scrub-successional bird community dynamics in young and mature longleaf pine-wiregrass savannahs, 803–814
Sawtooth National Forest: 607
Scandinavia: 728
Scaup, greater: 1362–1363
Scaup, lesser: 228, 384, 1362–1363
direct recovery rates of lesser scaup banded in northwest Minnesota: sources of heterogeneity, 389–395
Sceloporus jarrovi, see Lizard, spiny
Schaefer, James A., Alasdair M. Veitch, Fred H. Harrington, W. Kent Brown, John B. Theberge, and Stuart N. Luttich. Demography of decline of the Red Wine Mountains caribou herd, 580–587
Schaefferia glandulosa, see Yaupon, desert
Schaffer, Daniel, *see* Swindle, Keith A.
Schizachyrium scoparium, see Grass, little bluestem
Schmitz, Richard A., and William R. Clark. Survival of ring-necked pheasant hens during spring in relation to landscape features, 147–154
Schmitz, Richard A., *see* Clark, William R.
Schmutz, Joel A., and Craig R. Ely. Survival of greater white-fronted geese: effects of year, season, sex, and body condition, 1239–1249
Schwartz, Charles C., *see* Jacoby, Michael E.
Scirpus acutus, see Bulrush, hardstem
Scirpus spp., *see* Bulrush
Sciurus carolinensis, see Squirrel, gray
Sciurus niger, see Squirrel, fox
Sea star: 1066
Seaman, D. Erran, Joshua J. Millspaugh, Brian J. Kernohan, Gary C. Brundige, Kenneth J. Raedeke, and Robert A. Gitzen. Effects of sample size on kernal home range estimates, 739–747
Seamans, Mark E., *see* Peery, M. Zachariah
Seamans, Thomas W., *see* Blackwell, Bradley F.
Sedge: 287, 518, 854, 858, 898
Sedge, broom: 245

Sedge, elk: 347
Sedge, redroot flat: 1261–1268
Sedge, short-stalk: 336–337, 340–341
Sedge, tropical: 254
Sedgwick, James A., *see* Baker, Bruce W.
Seed production:
a new method to predict seed yield of moist-soil plants, 1269–1272
predicting seed yield of moist-soil plants, 1261–1268
Seiurus aurocapillus, see Ovenbird
Seiurus noveboracensis, see Thrush, northern water
Selasphorus rufus, see Hummingbird, rufous
Selenium:
selenium accumulation and effects on reproduction in captive American kestrels fed selenomethionine, 502–511
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
selenium in blood of predatory birds from Kesterson Reservoir and other areas in California, 1273–1281
Selenomethionine:
selenium accumulation and effects on reproduction in captive American kestrels fed selenomethionine, 502–511
Senecio serra, see Ragwort, golden
***Sequoia sempervirens*:** 45
Servheen, Chrostopher, *see* Jacoby, Michael E.
Serviceberry: 347, 1120
Sesbania spp., *see* Coffee, wild
Setaria virdis, see Grass, green bristle
Setophaga ruticilla, see Redstart, American
Sex allocation:
adaptive fetal sex allocation in elk: evidence and implications, 1109–1117
Sex ratio:
adaptive fetal sex allocation in elk: evidence and implications, 1109–1117
Shadscale: 774, 778
Shearwater, Newell's: 1000
Sheep: 1078
effect of preventive coyote hunting on sheep losses to coyote predation, 606–612
effects of summer sheep grazing on browse nutritive quality in autumn and winter, 346–354
relative vulnerability of coyotes to removal methods on a northern California ranch, 939–949
territoriality and breeding status of coyotes relative to sheep predation, 593–605
Sheep, Rocky Mountain bighorn: 1159
Shepherdia canadensis, see Buffalo berry
Shepperd, Wayne D., *see* Baker, Dan L.
Shideler, Susan E., *see* Stoops, Monica A.
Shoveler, northern: 382–389, 657, 1359–1369
Shrew: 444
Shrew, northern short-tailed: 1175, 1178
Shrew, ornate: 1278
Shrike, loggerhead: 807–808, 1273–1281
Shrimp, brine: 1209
Shumake, Stephen A., Ray T. Sterner, and Stanley E. Gaddis. Repellents to reduce cable gnawing by northern pocket gophers, 1344–1349
Sialia sialis, see Bluebird, eastern
Sidle, John G., *see* Kirsch, Eileen M.
Sigmodon hispidus, see Rat, hispid cotton
Silktassel: 456
Silver, Scott C., *see* Ostro, Linde E.
Silverling: 117
Simpson, Scott A., *see* Herkert, James R.
Simpson's diversity index: 38–39, 980
Simpson's eveness index: 980
Siskin, pine: 170–177
Sisymbrium altissimum, see Mustard, tumble
Sitta canadensis, see Nuthatch, red-breasted
Sitta carolinensis, see Nuthatch, white-breasted
Sitta pusilla, see Nuthatch, brown-headed
Siuslaw National Forest: 485
Skimmer, black:
use of dredged material substrates by nesting terns and black skimmers, 137–146
Skunk, striped: 370, 687, 985, 1070–1073, 1300
Slack:
slack in the configuration of habitat patches for northern bobwhites, 245–250
Smartweed: 359, 640
Smartweed, swamp: 1261–1268
Smilax spp., *see* Greenbrier
Smith, Adam, and Tomas Willebrand. Mortality causes and survival rates of hunted and unhunted willow grouse, 722–730
Smith, David A., *see* Williams, Byron K.
Smith, Winston P. Relations of small mammal populations to even-aged shelterwood systems: a comment, 1376–1380
Snag: 487, 1306–1319
Snail, lymnaeid: 362
Snake, black rat: 1286
Snake, brown tree: 298, 1009, 1015
Snake, garter: 445, 1175
Snake, gray rat: 224
Snake, Key ringneck: 258
Snake, northern ringneck: 1175
Snake River Birds of Prey National Conservation Area: 774–780
Snell, Howard L., *see* Gibbs, James P.
Snowberry: 346–354
Sodium cyanide: 939–949
Sodium monofluoroacetate: 1138–1151
Solenopsis invicta, see Ant, red fire
Solidago spp., *see* Goldenrod
Somateria mollissima, see Eider, common
Sonoran Desert: 77, 155
Sora: 122
Sorex ornatus, see Shrew, ornate
Sorex spp., *see* Shrew

Sorghastrum nutans, see Grass, Indian
South Carolina: 228, 283, 329, 803–814
South Dakota: 275, 386, 428–429, 470–483, 639–649, 662, 711–722
Soviet Union: 1090
Soybean: 956–963, 980
Sparrow, Bachman's: 803–814
Sparrow, Brewer's: 271
Sparrow, chipping: 807–809
Sparrow, field: 527, 530–535, 807
Sparrow, fox: 170–177
Sparrow, grasshopper: 530–538
Sparrow, Harris's: 228
Sparrow, house: 228, 1000, 1007, 1010, 1014–1015
Sparrow, Lincoln's: 170–177
Sparrow, sage: 271
Sparrow, savannah: 532
Sparrow, song: 188, 489–490
Sparrow, vesper: 532
Sparrow, white-crowned: 228, 488–491
Sparrow, white-throated: 170–177, 188
***Spartina bakeri*:** 1194
Spartina bakeri, see Grass, cord
Spartina bakeri, see Grass, sand cord
Spathodea campanulata, see Tree, African tulip
Spermophilus beechyi, see Squirrel, California ground
Spermophilus silosoma, see Squirrel, spotted ground
Spermophilus spp., *see* Squirrel, ground
Sphagnum spp., *see* Moss, peat
Spheniscus mendiculus, see Penguin, Galápagos
Sphyrapicus ruber, see Sapsucker, red-breasted
Sphyrapicus varius, see Sapsucker, yellow-bellied
Spirea betufolia, see Spirea, white
Spirea, white: 347
Spiza americana, see Dickcissel
Spizella breweri, see Sparrow, Brewer's
Spizella passerina, see Sparrow, chipping
Spizella pusilla, see Sparrow, field
Sporobolus spp., *see* Dropseed
Spruce, black: 168, 462, 468, 581, 1307
Spruce budworm: 462, 467
Spruce, Engelmann: 316
Spruce, Norway: 355, 812
Spruce, red: 181, 442, 461–466
Spruce, Sitka: 16, 21
Spruce, white: 168, 575, 581, 1307, 1377, 1382
Squirrel, California ground: 931, 1278
Squirrel, fox: 634–636
Squirrel, gray: 1173
Squirrel, ground: 788, 795, 965
Squirrel, northern flying:
movements of northern flying squirrels in different-aged forest stands of western Oregon, 291–297
Squirrel, San Joaquin antelope: 931
Squirrel, southern flying: 635, 825
effects of southern flying squirrels on nest success of red-cockaded woodpeckers, 538–545
Squirrel, spotted ground: 1067, 1071–1072, 1075, 1077
St. Johns National Wildlife Refuge: 116–125
Stanley, Thomas R., *see* Baker, Bruce W.
Starling, European: 228, 1007–1008, 1011, 1014, 1280
Statistical significance:
the insignificance of statistical significance testing, 763–772
Steenhof, Karen, *see* Kochert, Michael N.
Steffen, David E., *see* Pack, James C.
Stehn, Robert A., *see* Ward, David H.
Stephens, Scott E., *see* Blums, Peter
Sterna antillarum, see Tern, least
Sterna forsteri, see Tern, Forster's
Sterna nilotica, see Tern, gull-billed
Sterner, Ray T., *see* Shumake, Stephen A.
Stilt, black-necked: 1273
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
Stolley, Dorie S., John A. Bissonette, John A. Kadlec, and Dan Coster. Effects of saline drinking water on early gosling development, 990–996
Stoops, Monica A., Gary B. Anderson, Bill L. Lasley, and Susan E. Shideler. Use of fecal steroid metabolites to estimate the pregnancy rate of a free-ranging herd of tule elk, 561–569
Stratiomyids: 362
Strena hirundo, see Tern, common
Streptopelia chinensis, see Dove, spotted
Strix aluco, see Owl, tawny
Strix occidentalis caurina, see Owl, northern spotted
Strix occidentalis lucida, see Owl, Mexican spotted
Strix occidentalis occidentalis, see Owl, California spotted
Strix varia, see Owl, barred
Strychnine: 790
Sturnella magna, see Meadowlark, eastern
Sturnella neglecta, see Meadowlark, western
Sturnus vulgaris, see Starling, European
Sullivan, Druscilla S., *see* Sullivan, Thomas P. Sullivan, J. Daniel, Jr., and Ralph Mirarchi. Molt of after-hatching-year mourning doves, 223–231
Sullivan, Thomas P., Druscilla S. Sullivan, and Charlotte Kurta. Relations of small mammal populations to even-aged shelterwood systems: a reply, 1381–1389
Sulphur: 14–25
Sumner, Misty L., *see* Giuliano, William M.
Sunfish: 1166
Surnia ulula, see Owl, northern hawk
Survey, aerial: 472
Survey, broadcast:
efficacy of northern goshawk broadcast surveys in Washington state, 98–106
Survey technique:
dawn and morning goshawk courtship vocalizations as a method for detecting nest sites, 511–516

double-survey estimates of bald eagle populations in Oregon, 794–802
efficacy of northern goshawk broadcast surveys in Washington state, 98–106
influence of point count length and repeated visits on habitat model performance, 815–823
Survival, brood:
passive integrated transponders and patagial tag markers for northern bobwhite chicks, 162–166
wild turkey poult survival in southcentral Iowa, 199–203
Survival:
declining survival of ring-necked pheasant chicks in Illinois during the late 1900s, 705–710
direct recovery rates of lesser scaup banded in northwest Minnesota: sources of heterogeneity, 389–395
effect of Hurricane Andrew on survival and movements of white-tailed deer in the everglades, 872–879
effects of ear-tagging with radiotransmitters on survival of moose calves, 354–358
effects of fall hunting on wild turkey populations in Virginia and West Virginia, 964–975
effects of radiotransmitters on survival and reproductive success of gray partridge, 1044–1051
effects of violation of assumptions for survival analysis methods in radiotelemetry studies, 1369–1375
evaluation of a plasticine-filled leg band for day-old ducklings, 656–663
factors influencing wild turkey hen survival in southcentral Iowa, 731–738
macrohabitat effects on summer survival, movements, and clutch success of northern bobwhite in Kansas, 675–685
mortality causes and survival rates of hunted and unhunted willow grouse, 722–730
movements and survival of bald eagles banded in the Greater Yellowstone Ecosystem, 781–793
movements, survival, and mortality of black-tailed deer in the Klickitat basin of Washington, 861–871
mule deer survival in Colorado, Idaho, and Montana, 315–326
northern bobwhite chick mortality caused by red imported fire ants, 1291–1298
predation and survival of white-tailed deer fawns in northcentral New Brunswick, 574–579
rates and causes of grizzly bear mortality in the interior mountains of British Columbia, Alberta, Montana, Washington, and Idaho, 911–920
reproduction and survival of Rio Grande turkeys in Oregon, 204–210
survival and cause-specific mortality of adult bobcats in central Mississippi, 613–620
survival and cause-specific mortality of adult raccoons in central Mississippi, 880–888
survival and habitat use by ring-necked pheasants during two diparate winters in South Dakota, 711–722
survival and movements of translocated raccoons in northcentral Illinois, 278–286
survival and movements of orphaned white-tailed deer fawns in Texas, 570–574
survival of greater white-fronted geese: effects of year, season, sex, and body condition, 1239–1249
survival of postfledging mallards in northcentral Minnesota, 403–408
survivorship of a nonharvested raccoon population in south Texas, 889–894
Sus scrofa, see Pig, feral
Sustainability:
towards wildlife management in tropical forests, 1–13
Swallow, tree: 188
Swallow, violet-green: 489
Swan, Bewick's: 621
Swan, trumpter: 426
Swanson, David A., *see* Pack, James C.
Sweden: 354–358, 722–730
Sweitzer, Richard A., *see* Waithman, John D.
Swenson, Jon E., Kjell Wallin, Goran Ericsson, Goran Cederlund, and Finn Sandegren. Effects of ear-tagging with radiotransmitters on survival of moose calves, 354–558
Swindle, Keith A., William J. Ripple, E. Charles Meslow, and Daniel Schaffer. Old-forest distribution around spotted owl nests in the central Cascade Mountains, Oregon, 1212–1221
Sycamore: 451–459
Sycamore, American: 630–636
Sylvatic plague: 270–272
Sylvilagus audubonii, see Rabbit, desert cottontail *Sylvilagus floridanus, see* Rabbit, eastern cottontail
Sylvilagus nuttalli, see Rabbit, desert cottontail
Sylvilagus transitionalis, see, Rabbit, New England cottontail
Symphoricarpus spp., *see* Snowberry
Syzygium jambos, see Rose-apple
Sachycineta bicolcens, see Swallow, violet-green

T

Tachycineta bicolor, see Swallow, tree
Tadorna tadorna, see Duck, common shel
Tadorna tadornoides, see Duck, mountain
Tag, ear:
effects of ear-tagging with radiotransmitters on survival of moose calves, 354–358
Tag, patagial:
passive integrated transponders and patagial tag markers for northern bobwhite chicks, 162–166
Tag, web: 656–663
Taiwan: 1–13
Tallahaia Wildlife Management Area: 210–222, 261–269, 613–620, 880–888
Tamarind, wild: 874

Tamshiyacu-Tahuayo Community Reserve: 3–13
Tanager, scarlet: 184–188, 816–821
Tanager, summer: 807
Tanager, western: 488–490
Tapir, lowland: 3, 5
Tapirus terrestris, see Tapir, lowland
Taxidea taxus, see Badger
Tayassu pecari, see Peccary, white-lipped
Tayassu tajaeu, see Peccary, collared
Taylor, Curtis I., *see* Pack, James C.
Taylor, J. Scott, Kevin E. Church, and Donald H. Rusch. Microhabitat selection by nesting and brood-rearing northern bobwhite in Kansas, 686–694
———, and John R. Cary. Macrohabitat effects on summer survival, movements, and clutch success of northern bobwhite in Kansas, 675–685
Teal, blue-winged: 382–389, 418, 429, 647, 1359–1369
Teal, cinnamon: 1363–1364
Teal, green-winged: 384, 1359–1369
***Tectura* spp:** 17
Temperature: 1249–1260
influence of precipitation timing and summer temperatures on reproduction of Gambel's quail, 154–161
use of climatic data to model the presence of oestrid flies in caribou herds, 588–593
Tennessee: 283, 470–483, 815–823
Tern, common: 165
Tern, Forster's: 138
Tern, gull-billed:
use of dredged material substrates by nesting terns and black skimmers, 137–146
Tern, least:
status of the interior population of least tern, 470–483
use of dredged material substrates by nesting terns and black skimmers, 137–146
Tetrao tetrix, see Grouse, black
Texas: 228, 329, 451, 470–483, 539, 546, 551, 570–574, 824–832, 889–894, 995, 1009, 1066–1081, 1291–1305
Thalictrum fendler, see Rue, meadow
Thamnophis sirtalis, see Snake, garter
Theberge, John B., *see* Schaefer, James A.
Thermoregulation: 33, 855, 1128, 1147
Thief Lake Wildlife Management Area: 389–395
Thimbleberry: 346–354
Thistle: 347
Thistle, Russian: 1300
Thogmartin, Wayne E., and James E. Johnson. Reproduction in a declining population of wild turkeys in Arkansas, 1281–1290
Thome, Darrin M., Cynthia J. Zabel, and Lowell V. Diller. Forest stand characteristics and reproduction of northern spotted owls in managed north-coastal California forests, 44–59
Thomomys talpoides, see Gopher, northern pocket
Thompson, Ian D., *see* Jung, Thomas S.
Thompson, John D., *see* Mosillo, Maia
Thrasher, brown: 807
Thrasher, pearly-eyed: 242
Three-awn: 1067
Thrush, hermit: 170–177, 187–188
Thrush, northern water: 170–177, 188
Thrush, red-legged: 242
Thrush, Swainson's: 170–177, 184–188, 484–496
Thrush, varied: 484–496
Thrush, wood: 183–188, 184–188, 807
Thryomanes bewickii, see Wren, Bewick's
Thryothorus ludovicianus, see Wren, Carolina
Thuja accidentalis, see Cedar, eastern white
Thuja plicata, see Cedar, western red
Tichosurus caninus, see Possum, mountain brushtail
Tiger: 305
Timothy: 518, 575, 1328
Timothy, swamp: 359
Tipulids: 362
Tit, wren: 489–490
Titman, Rodger D., *see* Jung, Thomas S.
Titmouse, tufted: 807–809
Toad, American: 444, 448
Tongass National Forest: 1379
Torgos tracheliotus negevensis, see Vulture, negev lappet-faced
Tortoise, desert: 94
effects of drought on desert tortoise movement and activity, 1181–1192
Tortoise, Galápagos: 1060
Tortoise, Gopher:
burrow abandonment by gopher tortoises in slash pine plantations of the Conecuh National Forest, 26–35
Towhee, eastern: 807–809
Towhee, spotted: 488–492
Toxicity:
selenium accumulation and effects on reproduction in captive American kestrels fed selenomethionine, 502–511
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501
the efficacy of baits containing 1080 for control of brushtail possums, 1138–1151
Toxostoma rufum, see Thrasher, brown
Transect, line: 85–97, 232–244
Translocation:
evaluation of translocation for the threatened Florida scrub-jay, 833–842
further evaluation of the genetic consequences of translocations on southeastern white-tailed deer population, 327–334
real-time physiological responses of red deer to translocations, 1152–1162
survival and movements of translocated raccoons in northcentral Illinois, 278–286

Transponders, passive integrated:
passive integrated transponders and patagial tag markers for northern bobwhite chicks, 162–166
Trap, pitfall: 441–450
Trap, Tomahawk box: 881, 890
Tree, African tulip: 233
Tree, groundsel: 878
Tree, trumpet: 233
Trent-Rongstad method:
effects of violation of assumptions for survival analysis methods in radiotelemetry studies, 1369–1375
Trichechus manatus, see Manatee, Florida
Trichomonas gallinae, see Trichomoniasis
Trichomoniasis: 79–84
Trichosurus vulpecula, see Possum, brushtail
Trifolium spp., *see* Clover
Troglodytes aedon, see Wren, house
Troglodytes troglodytes, see Wren, winter
Trogon elegans, see Trogon, elegant
Trogon, elegant:
multiscaled habitat selection by elegant trogons in southeastern Arizona, 451–461
Trogon, rufous: 457
Trogon rufus, see Trogon, rufous
Tropidurus spp., *see* Lizard, lava
Trout, cutthroat: 788, 927
Trout, rainbow: 1016
Tsai, Kuenhi, Kenneth H. Pollock, and Cavell Brownie. Effects of violation of assumptions for survival analysis methods in radiotelemetry studies, 1369–1375
Tsuga canadensis, see Hemlock, eastern
Tsuga heterophylla, see Hemlock, western
Tsuga mertensia, see Hemlock, mountain
Turdus migratorius, see Robin, American
Turdus plumbeus, see Thrush, red-legged
Turkey, eastern wild: 163,165–166, 207–208
effects of fall hunting on wild turkey populations in Virginia and West Virginia, 964–975
factors influencing wild turkey hen survival in southcentral Iowa, 731–738
habitat use of eastern wild turkeys in central Mississippi, 210–222
reproduction in a declining population of wild turkeys in Arkansas, 1281–1290
wild turkey poult survival in southcentral Iowa, 199–203
Turkey, Merriam's: 208
Turkey, Rio Grande:
reproduction and survival of Rio Grande turkeys in Oregon, 204–210
Turnstone, ruddy: 144, 1000
Turtle, Blanding's: 858
Turtle, bog:
home range and habitat selection of bog turtles in southwestern Virginia, 853–860
Tympanuchus cupido pinnatus, see Chicken, greater prairie
Typha spp., *see* Cattail
Tyrannus forficatus, see Flycatcher, scissor-tailed
Tyrannus tyrannus, see Kingbird, eastern
Tyrannus verticalis, see Kingbird, western
Tyto alba, see Owl, barn

U

U. S. Department of Agriculture: 420, 422
U. S. Fish and Wildlife Service: 417–440, 498, 511, 517, 539–540, 553, 783–786, 833–834, 853, 951, 1240
U. S. Forest Service: 451, 607, 804–811, 816, 825, 881
Uinta National Forest: 607
Ulmus americana, see Elm, American
Ultrasonic detectors: 553–561
Umbellularia californica, see Bay, California
Unsworth, James W., David F. Pac, Gary C. White, and Richard M. Bartmann. Mule deer survival in Colorado, Idaho, and Montana, 315–326
Urban environment: 77–84, 930–938
Urine:
sampling and statistical analysis of snow-urine allantoin:cretinine ratios, 1118–1132
Urocyon cinereoargenteus, see Fox, gray
Ursus americanus, see Bear, black
USDA, *see* U. S. Department of Agriculture
USFWS, *see* U. S. Fish and Wildlife Service
Utah: 126–136, 270–277, 347, 606–612, 784, 990–996, 1078, 1327, 1349–1358

V

Vaccinium angustifolium, see Blueberry
Vaccinium spp., *see* Huckleberry
Vacine:
biosafety of parenteral *Brucella abortus* RB51 vaccine in bison calves, 950–955
Van Vuren, Dirk, *see* Waithman, John D.
Varley, John D., *see* Drewien, Roderick C.
Veery: 184–188
Veitch, Alasdair M., *see* Schaefer, James A.
Veratrum viride, see Indian poke
Verbascum thapsis, see Mullein
Vermivora celata, see Warbler, orange-crowned
Vermivora ruficapilla, see Warbler, Nashville
Vermont: 180–188
Vireo, blue-headed: 187–188
Vireo flavifrons, see Vireo, yellow-throated
Vireo gilvus, see Vireo, warbling
Vireo griseus, see Vireo, white-eyed
Vireo huttoni, see Vireo, Hutton's
Vireo, Hutton's: 489
Vireo olivaceus, see Vireo, red-eyed
Vireo, red-eyed: 188, 807, 816–821
Vireo solitarius, see Vireo, blue-headed
Vireo, solitary, *see* Vireo, blue-headed
Vireo, warbling: 484–496
Vireo, white-eyed: 807

Vireo, yellow-throated: 807
Virginia: 570, 811, 853–860, 964–975
Vocalization: 155
dawn and morning goshawk courtship vocalizations as a method for detecting nest sites, 511–516
efficacy of northern goshawk broadcast surveys in Washington state, 98–106
the effectiveness of tape playback in estimating black rail density, 116–125
Vole, California: 1278
Vole, northern red-backed: 1386
Vole, root: 745
Vole, southern redback: 1175, 1178
relations of small mammal populations to even-aged shelterwood systems: a comment, 1376–1380
relations of small mammal populations to even-aged shelterwood systems: a reply, 1381–1389
VORTEX: 251–260
Vulpes macrotis mutica, see Fox, San Joaquin kit
Vulpes velox, see Fox, swift
Vulpes vulpes, see Fox, red
Vultur gryphus, see Condor, Andean
Vulture, bearded: 112
Vulture, black: 113
Vulture, Egyptian:
supplemental feeding regimes for Egyptian vultures in the Negev Desert, Israel, 107–115
Vulture, Negev lappet-faced: 108
Vulture, turkey: 112, 1067, 1069

W

Waas, Joseph R., John R. Ingram, and Lindsay R. Matthews. Real-time physiological responses of red deer to translocations, 1152–1162
Wagner, Kimberly K., and Michael R. Conover. Effect of preventive coyote hunting on sheep losses to coyote predation, 606–612
Waithman, John D., Richard A. Sweitzer, Dirk Van Vuren, John D. Drew, Amy J. Brinkhaus, Ian A. Gardner, and Walter M. Boyce. Range expansion, population sizes, and management of wild pigs in California, 298–308
Wakkinen, Wayne L., *see* McLellan, Bruce N.
Walk, Jeff W., *see* Herkert, James R.
Wallin, Kjell, *see* Swenson, Jon E.
Walnut, black: 452
Walters, Jeffery R., *see* Zwicker, Susan M.
Warbler, black-and-white: 170–177, 807, 812
Warbler, black-throated blue: 183–188
Warbler, black-throated gray: 489
Warbler, black-throated green: 170–177, 183–188
Warbler, blackburnian: 184–188
Warbler, blackpoll: 170–177, 183–188
Warbler, Canada: 184–188
Warbler, chestnut-sided: 184–188, 816–821
Warbler, hermit: 484–496
Warbler, hooded: 807
Warbler, Kentucky: 807, 812
Warbler, MacGillivray's: 488–492
Warbler, magnolia: 170–177, 184–188
Warbler, mourning: 170–177, 188
Warbler, Nashville: 188
Warbler, orange-crowned: 488–492
Warbler, pine: 807–809
Warbler, prairie: 806–809
Warbler, Wilson's: 170–177, 484–496
Warbler, worm-eating: 816–821
Warbler, yellow: 170–177, 807
Warbler, yellow-rumped: 170–177, 188
Warbler, yellow-throated: 807
Ward, David H., Robert A. Stehn, Wallace P. Erickson, and Dirk V. Derksen. Response of fall-staging brant and Canada geese to aircraft overflights in southwestern Alaska, 373–381
Warner, Richard E., Philip C. Mankin, Larry M. David, and Stanley L. Etter. Declining survival of ring-necked pheasant chicks in Illinois during the late 1900s, 705–710
Warner, Richard E., *see* Mankin, Philip C.
Wasatch Cache National Forest: 607
Washington: 98–106, 292, 346, 784, 795, 911–920, 1103, 1162–1171
Water control structure: 426
Waterfowl: 358–368, 373–381, 382–389, 403–408, 621–663, 1299–1305
estimating statistical power to evaluate ongoing waterfowl population monitoring, 1359–1369
evaluation of waterfowl conservation under the North American Waterfowl Management Plan, 417–440
water and sediment characterics associated with avian botulism outbreaks in wetland, 1249–1260
Watson, James W., David W. Hays, and D. John Pierce. Efficacy of northern goshawk broadcast surveys in Washington state, 98–106
Watts, E. Jane, *see* Flemming, Stephen P.
Waxwing, cedar: 187–188, 807
Weasel: 202, 292
Weasel, long-tailed: 275
Weeks, Harmon P. Jr., *see* Frawley, Brian J.
Weerakkody, Govinda, *see* Gray, Matthew J.
Welder Wildlife Refuge: 889–894
West Virginia: 964–975
Westemeier, Ronald L., *see* Herkert, James R.
Wetland: 417–440, 853–860, 898, 1162–1171, 1261–1281
environmental characteristics associated with the occurrence of avian botulism in wetlands of a northern California refuge, 358–368
resource variability and conservation of American flamingos in coastal wetlands of Yucatán, Mexico, 1201–1212
selenium concentrations in eggs of American avocets and black-necked stilts at an evaporation basin and freshwater wetland in California, 497–501

water and sediment characterics associated with avian botulism outbreaks in wetland, 1249–1260
Whale, right: 1352
Wheat: 956–963
Whip-poor-will: 807
Whitaker, Darroch M., and William A. Montevecchi. Breeding bird assemblages inhabiting riparian buffer strips in Newfoundland, Canada, 167–179
White, Gary C., *see* Otis, David L.
———, *see* Unsworth, James W.
White Mountain National Forest: 185
Whitebrush: 571
Whitefish, mountain: 788
Whitlaw, Heather A., *see* Ballard, Warren B.
Wigeon, American: 384, 429–430, 1363–1364
Wigeon, Eurasian: 1231
Willamette National Forest: 292, 1212–1221
Willebrand, Tomas, *see* Smith, Adam
Williams, Byron K., Mark D. Koneff, and David A. Smith. Evaluation of waterfowl conservation under the North American Waterfowl Management Plan, 417–440—Invited paper
Willow: 581, 630–636, 640, 877, 1321
Willow, Arctic: 287
Willow, Gooding's: 457
Willow, Scouler: 346–354
Wilson, Barry W., *see* Santolo, Gary M.
Wilson, Tim S., *see* Chamberlain, Michael J.
Wilsonia canadensis, *see* Warbler, Canada
Wilsonia citrina, *see* Warbler, hooded
Wilsonia pusilla, *see* Warbler, Wilson's
Winter: 346–354, 1087–1088, 1130
population energetics of northern pintails wintering in the Sacramento Valley, California, 1222–1238
sampling and statistical analysis of snow-urine allantoin:cretinine ratios, 1118–1132
winter prey selection by wolves and cougars in and near Glacier National Park, Montana, 901–910
Winter, Brian, *see* Merrill, Michael D.
Winterfat: 86, 774
Wisconsin: 329, 1103, 1287
Wolf: 109, 111
Wolf, Ethiopian: 936
Wolf, gray: 305, 335–336, 338, 341, 343, 574–579, 580–587, 603, 935, 1066
characteristics of dispersal in a colonizing wolf population in the central Rocky Mountains, 1094–1108
power analysis of wolf-moose functional responses, 396–402
species-specific population dynamics of cervids in a multipredator ecosystem, 1082–1093
winter prey selection by wolves and cougars in and near Glacier National Park, Montana, 901–910
Wolverine: 1104
Woodard, Allan R., *see* Rice, Kenneth G.
Woodpecker, black: 466
Woodpecker, black-bellied: 170–177
Woodpecker, downy: 170–177, 187–188, 489, 807
Woodpecker, European white-backed: 467
Woodpecker, hairy: 170–177, 187–188, 488–492, 807
Woodpecker, pileated: 188, 490, 630–636
characteristics of foraging trees selected by pileated woodpeckers in New Brunswick, 361–469
Woodpecker, red-bellied: 807
Woodpecker, red-cockaded: 467
effects of southern flying squirrels on nest success of red-cockaded woodpeckers, 538–545
estimation of habitat selection for central-place foraging animals, 1028–1038
reintroduction reintroduction and postrelease movements of red-cockaded woodpecker groups in eastern Texas, 824–832
scrub-successional bird community dynamics in young and mature longleaf pine-wiregrass savannahs, 803–814
selection of pines for foraging by red-cockaded woodpeckers, 843–852
Woodpecker, red-headed: 807
Woods, John G., *see* McLellan, Bruce N.
Wren, Bewick's: 489–490
Wren, Carolina: 807–809
Wren, house: 488–492, 633
Wren, winter: 170–177, 188, 484–496
Wyoming: 85–97, 126–136, 270–277, 369, 470–483, 594, 602, 606–612, 782–790, 921–929, 950–955, 1118–1132, 1327

X

Xanthium strumarium, *see* Cocklebur
Xylorhiza glabriuscula, *see* Aster, woody

Y

Yakama Indian Reservation: 861–871
Yamamoto, Julie T., *see* Santolo, Gary M.
Yaupon, desert: 571
Yellowstone National Park: 603, 901, 950–955, 1106, 1119–1132, 1327
brucellosis in Yellowstone National Park bison: quantitative serology and infection, 1132–1137
movements and survival of bald eagles banded in the Greater Yellowstone Ecosystem, 781–793
seasonal movements of sandhill cranes radiomarked in Yellowstone National Park and Jackson Hole, Wyoming, 126–136
Yellowthroat, common: 187–188, 489, 530–538, 807
Yersinia taxus, *see* Sylvatic plague
Yetter, Aaron P., Stephen P. Havera, and Christopher S. Hine. Natural-cavity use by nesting wood ducks in Illinois, 630–638
Young, Steven J., *see* Ballard, Warren B.
Young, Truman P., *see* Ostro, Linde E.
Yucatán: 1201–1212, 1231
Yucca: 456, 1067
Yucca spp., *see* Yucca

Z

Zabel, Cynthia J., *see* Thome, Darrin M.
Zaiglin, Robert E., *see* Giuliano, William M.
Zebra: 1158
Zenaida aurita, see Dove, Zenaida
Zenaida macroura, see Dove, mourning
Zendaida asiatica, see Dove, white-winged
Zimmerman, John L., *see* Hughes, John P.
Zonotrichia albicollis, see Sparrow, white-throated
Zonotrichia leucophyrs, see Sparrow, white-crowned
Zonotrichia querula, see Sparrow, Harris's
Zostera marina, see Grass, eel
Zwicker, Susan M., and Jeffery R. Walters. Selection of pines for foraging by red-cockaded woodpeckers, 843–852